

Ontario Respiratory Virus Bulletin | 2013-2014

SURVEILLANCE WEEK 6 (February 2, 2014 – February 8, 2014)

This issue of the Ontario Respiratory Virus Bulletin provides information on the surveillance period from February 2 to February 8, 2014 (Week 6). Unless otherwise specified, **data presented in this issue of the bulletin are for Week 6** and data extraction occurred on Wednesday, February 12, 2014. Weekly production of this expanded version of the bulletin will continue throughout the influenza season until May 2014, and then an abbreviated version will be published on alternate weeks.

Assessment of Influenza Activity in Ontario

Measure of activity	Assessment of measure compared to the previous week (Lower, Similar, Higher)	Reasoning behind assessment
Laboratory confirmed influenza cases ^{1,2}	Lower	There were 385 influenza cases (332 influenza A and 53 influenza B) reported ¹ in week 6 (Table 1), which was less than the number of influenza cases reported in week 5 (472 influenza A and 29 influenza B). Percent positivity ² of laboratory tests for all influenza types in Ontario was 16.5% for week 6 (Table 3), which was lower than percent positivity in week 5 (17.7%).
Influenza outbreaks ³	Higher	There were six new influenza outbreaks reported in week 6: three influenza A and three influenza B; this was higher than the three influenza outbreaks reported in week 5: two influenza A and one influenza B (Table 4).
Influenza activity levels reported by health units	Similar	Twenty health units reported 'sporadic', fourteen reported 'localized', and one reported 'widespread' levels of influenza activity in week 6 (Figure 6). In week 5, eighteen health units reported 'sporadic', sixteen reported 'localized', and one reported 'widespread' levels of influenza activity.
OVERALL ASSESSMENT	Similar	Overall, the indicators show that influenza activity for week 6 was similar compared to week 5.
Measures of severity for influenza		
Hospitalizations among lab confirmed cases of influenza ¹	Lower	There were 154 hospitalized cases reported in week 6, which was lower than the 179 hospitalized cases reported in week 5 (Table 2).
Deaths among lab confirmed cases of influenza ¹	Lower	There were 12 deaths reported in week 6, which was lower than the 13 deaths reported in week 5 (Table 2).


¹ The number of **reported** cases needs to be interpreted with caution because there is a lag between the onset of illness, hospital admission or death and when this information is reported to public health. ² Positivity among specimens submitted for testing to Ontario laboratories reporting to the Centre for Immunization and Respiratory Infectious Diseases (CIRID). ³ The number of new outbreaks reported for the current week is based on the onset date of illness for the first case in an outbreak.

Summary of current and season-to-date respiratory virus activity in Ontario

- Influenza activity in the current surveillance week of the 2013-2014 season was similar compared to the previous week. Influenza remains the dominant circulating respiratory virus. Based on laboratory findings¹, influenza A is currently the predominant circulating influenza type and H1N1pdm09 is the predominant circulating subtype.
 - Influenza percent positivity overall for the season continues to decline and has been declining since week 2. Influenza A percent positivity was 14.1% in week 6, which was slightly lower compared to week 5 (16.8%); however, influenza B percent positivity was 2.4% in week 6, which was higher compared to week 5 (0.9%).
 - A total of 5033 laboratory-confirmed influenza cases has been reported for the 2013-2014 surveillance season to date. Among these cases, the majority (96.3%, 4845/5033) were influenza A (Table 1).
 - In the season to date, 1730 hospitalizations and 104 deaths have been reported among laboratory-confirmed influenza cases (Table 2). The majority of hospitalizations have occurred among influenza A cases (97.2%, 1681/1730).
- Among all other circulating respiratory viruses in week 6 (Table 3):
 - Respiratory syncytial virus had the highest percent positivity (11.3%), which was slightly higher compared to the previous week (9.9%).
 - Coronavirus had the second highest percent positivity (8.1%), which was lower compared to the previous week (13.0%).

¹ Based on percent positivity among specimens submitted for testing to laboratories reporting to the Public Health Agency of Canada's Centre for Immunization and Respiratory Infectious Diseases (CIRID).

Figure 1. Number of reported confirmed cases of **influenza A** by surveillance week: Ontario, September 1, 2008 to February 8, 2014



Source: Ontario Ministry of Health and Long-Term Care, integrated Public Health Information System (iPHIS) database, extracted by Public Health Ontario [2014/02/12].


Notes:

Interpret most recent case counts for the current season with caution due to reporting lags.

*Case counts above are based on 'Reported Date', which was the date the health unit was notified of the case; however, cases are assigned to a particular surveillance week based on the episode date entered in iPHIS for the case. Episode date for a case corresponds to the earliest date on record for the case according to the iPHIS hierarchy (Symptom Date > Clinical Diagnosis Date > Specimen Collection Date > Lab Test Date > Reported Date). Cases are excluded from these counts if the episode date is based on reported date, and the reported date occurs after the latest surveillance week.

Cases with an episode date after week 15 in the 2008-2009 season, as well as all cases from the 2009-2010 influenza season, were excluded from this graph based on the assumption that the influenza burden during the pandemic period is not comparable to a regular influenza season. Week 53 of the 2008-2009 season has been removed for continuity in displaying trends.

Figure 2. Number of reported confirmed cases of **influenza B** by surveillance week: Ontario, September 1, 2009 to February 8, 2014



Source: Ontario Ministry of Health and Long-Term Care, integrated Public Health Information System (iPHIS) database, extracted by Public Health Ontario [2014/02/12].

Notes:

Interpret most recent case counts for the current season with caution due to reporting lags.

*Case counts above are based on 'Reported Date', which was the date the health unit was notified of the case; however, cases are assigned to a particular reporting week based on the episode date entered in iPHIS for the case. Episode date for a case corresponds to the earliest date on record for the case according to the iPHIS hierarchy (Symptom Date > Clinical Diagnosis Date > Specimen Collection Date > Lab Test Date > Reported Date). Cases are excluded from these counts if the episode date is based on reported date, and the reported date occurs after the latest surveillance week.

Table 1. Number of reported confirmed influenza cases by health unit and health region: Ontario, surveillance week 6 and cumulative total (in parentheses) for the 2013-2014 season

Health Unit and Region	Influenza A				Influenza A & B	Influenza B	TOTAL
	(H1N1) pdm09	No subtype available	Other *	H3			
Northwestern	0 (4)	3 (41)	0 (0)	0 (0)	0 (0)	0 (1)	3 (46)
Thunder Bay District	1 (41)	1 (71)	0 (0)	0 (0)	0 (0)	0 (1)	2 (113)
TOTAL NORTH WEST	1 (45)	4 (112)	0 (0)	0 (0)	0 (0)	0 (2)	5 (159)
Algoma	0 (24)	1 (44)	0 (1)	0 (0)	0 (0)	0 (0)	1 (69)
North Bay Parry Sound District	0 (35)	5 (157)	0 (0)	0 (0)	0 (0)	0 (6)	5 (198)
Porcupine	1 (8)	7 (32)	0 (0)	0 (0)	0 (0)	0 (1)	8 (41)
Sudbury & District	1 (19)	4 (100)	0 (0)	0 (0)	0 (0)	0 (1)	5 (120)
Timiskaming	0 (3)	2 (6)	0 (0)	0 (0)	0 (0)	0 (0)	2 (9)
TOTAL NORTH EAST	2 (89)	19 (339)	0 (1)	0 (0)	0 (0)	0 (8)	21 (437)
City of Ottawa	1 (41)	15 (184)	0 (0)	0 (9)	0 (0)	1 (4)	17 (238)
Eastern Ontario	0 (20)	1 (41)	0 (1)	0 (9)	0 (0)	0 (1)	1 (72)
Hastings & Prince Edward Counties	0 (7)	3 (10)	0 (0)	0 (0)	0 (0)	1 (1)	4 (18)
Kingston, Frontenac, Lennox & Addington	0 (45)	10 (49)	0 (0)	0 (21)	0 (0)	1 (7)	11 (122)
Leeds, Grenville And Lanark District	0 (26)	4 (30)	0 (0)	0 (0)	0 (0)	2 (5)	6 (61)
Renfrew County And District	0 (10)	2 (32)	0 (0)	0 (4)	0 (0)	0 (0)	2 (46)
TOTAL EASTERN	1 (149)	35 (346)	0 (1)	0 (43)	0 (0)	5 (18)	41 (557)
Durham Region	0 (54)	17 (106)	0 (0)	0 (5)	0 (0)	1 (1)	18 (166)
Haliburton, Kawartha, Pine Ridge	1 (26)	4 (52)	0 (0)	0 (2)	0 (0)	0 (1)	5 (81)
Peel Region	8 (184)	26 (260)	0 (0)	3 (23)	0 (0)	12 (32)	49 (499)
Peterborough County-City	6 (29)	2 (22)	0 (0)	0 (0)	0 (0)	0 (1)	8 (52)
Simcoe Muskoka District	4 (103)	7 (114)	0 (0)	0 (1)	0 (0)	1 (5)	12 (223)
York Region	2 (102)	20 (135)	0 (2)	0 (6)	0 (0)	7 (16)	29 (261)
TOTAL CENTRAL EAST	21 (498)	76 (689)	0 (2)	3 (37)	0 (0)	21 (56)	121 (1282)
Toronto	25 (607)	48 (315)	0 (0)	2 (55)	0 (0)	14 (55)	89 (1032)
TOTAL TORONTO	25 (607)	48 (315)	0 (0)	2 (55)	0 (0)	14 (55)	89 (1032)
Chatham-Kent	1 (28)	0 (34)	0 (0)	0 (0)	0 (1)	2 (3)	3 (66)
Elgin-St. Thomas	1 (21)	0 (19)	0 (0)	0 (0)	0 (0)	0 (0)	1 (40)
Grey Bruce	0 (20)	1 (57)	0 (0)	0 (0)	0 (0)	0 (0)	1 (77)
Huron County	0 (8)	3 (11)	0 (0)	0 (0)	0 (0)	0 (0)	3 (19)
Lambton County	1 (22)	3 (17)	0 (0)	0 (0)	0 (0)	0 (1)	4 (40)
Middlesex-London	1 (31)	6 (133)	0 (1)	0 (0)	0 (0)	4 (7)	11 (172)
Oxford County	0 (8)	0 (15)	0 (0)	0 (1)	0 (0)	0 (1)	0 (25)
Perth District	0 (14)	1 (15)	0 (0)	1 (5)	0 (0)	0 (2)	2 (36)
Windsor-Essex County	2 (110)	1 (65)	0 (0)	0 (0)	0 (0)	0 (0)	3 (175)
TOTAL SOUTH WEST	6 (262)	15 (366)	0 (1)	1 (6)	0 (1)	6 (14)	28 (650)
Brant County	1 (12)	5 (32)	0 (0)	0 (1)	0 (0)	0 (1)	6 (46)
City Of Hamilton	0 (15)	20 (211)	0 (0)	1 (3)	0 (0)	2 (20)	23 (249)
Haldimand-Norfolk	0 (9)	2 (24)	0 (0)	0 (4)	0 (0)	0 (1)	2 (38)
Halton Region	1 (42)	12 (118)	0 (0)	0 (4)	0 (0)	3 (6)	16 (170)
Niagara Region	1 (49)	11 (88)	0 (0)	0 (2)	0 (0)	1 (3)	13 (142)
Waterloo Region	2 (29)	14 (135)	0 (0)	0 (4)	0 (0)	1 (4)	17 (172)
Wellington-Dufferin-Guelph	0 (40)	3 (58)	0 (0)	0 (1)	0 (0)	0 (0)	3 (99)
TOTAL CENTRAL WEST	5 (196)	67 (666)	0 (0)	1 (19)	0 (0)	7 (35)	80 (916)
TOTAL ONTARIO	61 (1846)	264 (2833)	0 (5)	7 (160)	0 (1)	53 (188)	385 (5033)

Source: Ontario Ministry of Health and Long-Term Care, integrated Public Health Information System (iPHIS) database, extracted by Public Health Ontario [2014/02/12].

Notes: Case counts above are based on 'Reported Date', which was the date the health unit was notified of the case. The cumulative count (in parentheses) includes laboratory-confirmed cases with a 'Reported Date' between September 1, 2013 and the last day of the current surveillance week. * The 'Other' influenza A category includes indeterminate, untypeable and other subtypes.

Table 2. Hospitalizations and deaths among confirmed influenza cases by age group: Ontario, week 6 and cumulative for the 2013-2014 influenza season


Age Group	HOSPITALIZATIONS		DEATHS	
	Weekly Count	Cumulative count (rate per 100,000)	Weekly count	Cumulative count (rate per 100,000)
<1	12	134 (94.7)	0	1 (0.7)
1 – 4	14	208 (36.1)	0	2 (0.3)
5 – 14	12	74 (5)	0	2 (0.1)
15 – 24	1	49 (2.7)	0	2 (0.1)
25 – 44	23	223 (6)	1	9 (0.2)
45 – 64	47	514 (13.5)	8	45 (1.2)
65+	45	528 (26.7)	3	43 (2.2)
Total	154	1730 (12.8)	12	104 (0.8)

Sources:

Case data: Ontario Ministry of Health and Long-Term Care (MOHLTC), integrated Public Health Information System (iPHIS) database, extracted by Public Health Ontario [2014/02/12].

Population data: Population Estimates 1986-2012, Ontario Ministry of Health and Long-Term Care, IntelliHEALTH ONTARIO, extracted by Public Health Ontario [2013/09/16]

Figure 3. Percentage of respiratory viral pathogens (influenza A, influenza B, respiratory syncytial virus, and parainfluenza virus) detected among specimens tested by all methods: Ontario, February 10, 2013 to February 8, 2014



Source: These data have been obtained from the Public Health Agency of Canada's (PHAC) Centre for Immunization and Respiratory Infectious Diseases (CIRID) respiratory virus detection tables as of February 12, 2014; they are based on data submitted to PHAC from 16 laboratories in Ontario.


Notes:

The numbers reported in this figure represent results submitted to the CIRID by 16 participating laboratories in Ontario, including 11 Public Health Ontario Laboratories (PHOLs) and five hospital-based laboratories.

Interpret results for week 51 with caution as only four laboratories in Ontario, three of which were hospital-based laboratories, submitted results to the CIRID. Results above are assigned to a particular surveillance week based on when test results are reported to PHAC; these data are not updated when results are submitted late for previous surveillance weeks. These data represent the number of specimens tested, which may not necessarily correspond with the number of patients as more than one specimen may have been submitted per patient.

Cumulative numbers for the season to date are also available through FluWatch: <http://www.phac-aspc.gc.ca/fluwatch/>

Figure 4. Percentage of respiratory viral pathogens (adenovirus, human metapneumovirus, rhinovirus and coronavirus) detected among specimens tested by all methods: Ontario, February 10, 2013 to February 8, 2014



Source: These data have been obtained from the Public Health Agency of Canada's (PHAC) Centre for Immunization and Respiratory Infectious Diseases (CIRID) respiratory virus detection tables as of February 12, 2014; they are based on data submitted to PHAC from 16 laboratories in Ontario.

Notes:

The numbers reported in this figure represent results submitted to the CIRID by 16 participating laboratories in Ontario, including 11 Public Health Ontario Laboratories and five hospital-based laboratories.

Interpret results for week 51 with caution as only four laboratories in Ontario, three of which were hospital-based laboratories, submitted results to the CIRID.

Results above are assigned to a particular surveillance week based on when test results are reported to PHAC; these data are not updated when results are submitted late for previous surveillance weeks. These data represent the number of specimens tested, which may not necessarily correspond with the number of patients as more than one specimen may have been submitted per patient.

Cumulative numbers for the season to date are also available through FluWatch: <http://www.phac-aspc.gc.ca/fluwatch/>

Table 3. Number and percent positivity of respiratory specimens tested by all methods for influenza and other respiratory viruses: Ontario, week 6 and cumulative for the 2013-2014 season

Detected viruses	Current Surveillance Week			Cumulative for Season*		
	Number positive	Number tested	Percent positive	Number positive	Number tested	Percent positive
Influenza (All)	416	2,522	16.5%	3,645	25,215	14.5%
Influenza A	356	2,522	14.1%	3,572	25,215	14.2%
Influenza B	60	2,522	2.4%	73	25,215	0.3%
Parainfluenza virus	32	2,203	1.5%	621	20,878	3.0%
Adenovirus	19	2,203	0.9%	172	20,878	0.8%
Respiratory syncytial virus	283	2,494	11.3%	1,413	23,482	6.0%
Rhinovirus	19	688	2.8%	914	8,331	11.0%
Human metapneumovirus	52	2,150	2.4%	192	19,943	1.0%
Coronavirus	31	384	8.1%	131	5,229	2.5%

Source: These data have been obtained from the Public Health Agency of Canada's (PHAC) Centre for Immunization and Respiratory Infectious Diseases (CIRID) respiratory virus detection tables as of February 12, 2014; they are based on data submitted to PHAC from 16 participating laboratories in Ontario.

Notes:

*Cumulative numbers include counts from August 25, 2013 (surveillance week 35) because PHAC's CIRID data include week 35 as part of the 2013-2014 season.

The data in this table are based on the date on which test results are reported. Cumulative counts reported include late reports from previous weeks; however, the current week will not include data from laboratories that submit late reports. Therefore, weekly counts from previous weeks may not equal the cumulative count.

These data represent the number of specimens tested, which may not necessarily correspond with the number of patients as more than one specimen may have been submitted per patient.

Cumulative numbers for the season to date are also available through FluWatch: <http://www.phac-aspc.gc.ca/fluwatch/>

Table 4. Institutional respiratory infection outbreaks: Ontario, week 6 and total for the 2013-2014 season to date¹

Institutional Respiratory Infection Outbreaks		
New outbreaks (Week 6) ²	Number of outbreaks	Percentage (%) of total
Influenza A	3	37.5
Influenza B	3	37.5
Influenza A and B	0	0.0
Enterovirus/rhinovirus	0	0.0
Parainfluenza (All types)	0	0.0
Respiratory Syncytial Virus (RSV)	0	0.0
Combined outbreaks ³	0	0.0
Other organisms ⁴	0	0.0
No organism identified	2	25.0
TOTAL	8	100.0
Cumulative outbreaks (season to date) ¹	N	
Influenza A	65	16.6
Influenza B	10	2.6
Influenza A and B	0	0.0
Enterovirus/rhinovirus	114	29.1
Parainfluenza (All types)	30	7.7
Respiratory Syncytial Virus (RSV)	31	7.9
Combined outbreaks ³	21	5.4
Other organisms ⁴	38	9.7
No organism identified	83	21.2
TOTAL	392	100.0
Types of institutions for cumulative outbreaks		
Long-Term Care Homes	248	63.3
Hospitals	31	7.9
Retirement Homes	53	13.5
Other ⁵	1	0.3
Unknown	59	15.1
TOTAL	392	100.0

Source: Ontario Ministry of Health and Long-Term Care, integrated Public Health Information System (iPHIS) database, extracted by Public Health Ontario [2014/02/12].

Notes:

¹ Season to date includes all outbreaks in which the date of onset of illness for the first case associated with the outbreak occurred between September 1, 2013 (Week 36) and the current reporting week, as well as outbreaks in which the date of onset of illness for the first case is missing.


² New outbreaks are those in which the date of onset of illness in the first case occurred and was recorded in iPHIS in the current surveillance period; i.e. February 2 – February 8, 2014 (Week 6).

³ Combined outbreaks include outbreaks in which more than one non-influenza organism has been identified (e.g. RSV, parainfluenza, rhinovirus, etc.).

⁴ Other organisms include outbreaks involving other aetiological agents, such as human metapneumovirus, adenovirus, and coronavirus.

⁵ Other types of institutions include: correctional facilities, group homes, shelters, and facilities operating under the *Developmental Services Act*. Note that school-based respiratory outbreaks are not captured in this table.

Figure 5. Institutional respiratory infection outbreaks by week of illness onset in the first case: Ontario, February 10, 2013 to February 8, 2014




Source: Ontario Ministry of Health and Long-Term Care, integrated Public Health Information System (iPHIS) database, extracted by Public Health Ontario [2014/02/12].

Notes:

Institutional respiratory infection outbreaks for which the date of onset of illness for the first case is missing are excluded in this figure. However, these outbreaks are counted in the cumulative outbreaks section of Table 4.

Figure 6. Influenza activity levels reported by public health units: Ontario, February 2 to February 8, 2014 (Week 6)




Source: Public Health Ontario [Provincial Influenza Activity Report (Appendix C) Database]

Notes:

Influenza activity levels are assigned by local public health units and reported to Public Health Ontario by the Tuesday following the end of each surveillance week at 4:00 p.m. Activity levels are assigned based on laboratory confirmations, ILI reports from various sources, and laboratory-confirmed institutional respiratory infection outbreaks. Please click [here](#) for [detailed definitions for the 2013-2014 season](#).

Activity levels reported for a particular surveillance week may not necessarily correspond to the number of new outbreaks reported in the same week in Table 3 because ongoing outbreaks from previous weeks, as well as laboratory confirmed outbreaks in schools, may be included in the assessment of the activity level.

Figure 7. Influenza-like-illness (ILI) consultation rate (per 1,000 patient visits) reported by sentinels by surveillance week compared to historical average (2007/2008 - 2012/2013 seasons): Ontario, February 10, 2013 to February 8, 2014



Source: Public Health Agency of Canada, FluWatch ILI sentinel surveillance reporting

Notes:

Sentinel information is reported to the Public Health Agency of Canada; 57 sentinel sites reported in week 6. Due to the small number of sentinel sites reporting each week, readers are advised to consider overall trends and place less emphasis on week-to-week fluctuations.

Ontario average ILI rates include data from the 2007-2008 to the 2012-2013 seasons, excluding ILI rates during the time period of the influenza pandemic in Ontario (April 2009 to January 2010) due to the incomparability of those rates to a regular influenza season.

Table 5. Influenza-like illness (ILI) consultation rate (per 1,000 patient visits) by age group as reported by sentinel sites: Ontario, week 6

Age Category	0 – 4	5 – 19	20 – 64	65 +	Unknown	TOTAL
Number of Patients with ILI	4	11	38	5	0	58
Number of Total Consultations	96	122	753	345	0	1316
<i>ILI Consultation Rate (per 1,000 patient visits)</i>	41.7	90.2	50.5	14.5	0	44.1

Source: Public Health Agency of Canada, FluWatch ILI sentinel surveillance reporting

Notes:

Sentinel information is reported to the Public Health Agency of Canada; 57 sentinel sites reported in week 6. ILI consultation rates presented in Table 5 should be interpreted with caution, as small numbers of sentinel sites reporting can result in unstable ILI rates.

APPENDIX**Table I.** Strain characterization completed on influenza isolates at the National Microbiology Laboratory: Ontario and Canada, September 1, 2013 to February 13, 2014

Influenza strains	Ontario	Canada
Influenza A (H3N2) A/Texas/50/2012-like	33	48
Influenza A (H1N1)pdm09 A/California/07/2009-like	259	755
Influenza B B/Brisbane/60/2008-like	1	9
Influenza B B/Massachusetts/02/2012-like	32	82

Source: Influenza and Respiratory Viruses Section, National Microbiology Laboratory (NML).

Notes:

The influenza strains recommended by the World Health Organization (WHO) for the Northern Hemisphere vaccine for the [2013-2014 season](#) are: A/California/7/2009-like (A H1N1pdm09-like virus); A/Texas/50/2012-like virus; and B/Massachusetts/02/2012-like virus (Yamagata lineage).


Table II. Amantadine, oseltamivir and zanamivir susceptibility assays completed on influenza isolates at the National Microbiology Laboratory: Ontario and Canada, September 1, 2013 to February 13, 2014

Influenza strains	Amantadine				Oseltamivir				Zanamivir			
	ONTARIO		CANADA		ONTARIO		CANADA		ONTARIO		CANADA	
	R	S	R	S	R	S	R	S	R	S	R	S
Influenza A (H3N2)	37	0	53	0	0	28	0	41	0	28	0	41
Influenza A (H1N1)pdm09	202	0	496	0	0	192	0	472	0	189	0	463
Influenza B	NA	NA	NA	NA	0	19	0	48	0	19	0	48

(R = Resistant, S = Susceptible, NA = Not Applicable)

Source: Influenza and Respiratory Viruses Section, National Microbiology Laboratory (NML).

Figure I. Total number of influenza tests performed and percent of positive tests by report week: Ontario, February 3, 2013 to February 8, 2014



Source: These data have been obtained from the Public Health Agency of Canada's (PHAC) Centre for Immunization and Respiratory Infectious Diseases (CIRID) respiratory virus detection tables as of February 12, 2014; they are based on data submitted to PHAC from 16 laboratories in Ontario.

Notes:

The numbers reported in this figure represent results submitted to the CIRID by 16 participating laboratories in Ontario, including 11 Public Health Ontario Laboratories and five hospital-based laboratories.

Interpret results for week 51 with caution as only four laboratories in Ontario, three of which were hospital-based laboratories, submitted results to the CIRID.

Results above are assigned to a particular surveillance week based on when test results are reported to PHAC; these data are not updated when results are submitted late for previous surveillance weeks.

These data represent the number of specimens tested, which may not necessarily correspond with the number of patients as more than one specimen may have been submitted per patient.

Cumulative numbers for the season to date are also available through FluWatch: <http://www.phac-aspc.gc.ca/fluwatch/>