

Ontario Respiratory Pathogen Bulletin | 2015-2016

SURVEILLANCE WEEK 43 (October 25, 2015 – October 31, 2015)

This issue of the Ontario Respiratory Pathogen Bulletin provides information on the surveillance period from October 25 to October 31, 2015 (Week 43). Unless otherwise specified, **data presented in this issue of the bulletin are for Week 43** and data extraction occurred on Wednesday, November 4, 2015.

Assessment of Influenza Activity in Ontario

Measure of activity	Assessment of measure compared to the previous week (Lower, Similar, Higher)	Reasoning behind assessment
Laboratory-confirmed influenza cases ¹	Similar	There were 9 influenza cases (5 influenza A and 4 influenza B) reported in Week 43, which was similar to the 8 influenza cases reported in Week 42 (all influenza A).
Percent positivity for influenza ²	Similar	Percent positivity of laboratory tests for all influenza types in Week 43 was 0.9%, which was similar to the percent positivity in Week 42 (0.8%). In Week 43, percent positivity was 0.6% for influenza A and 0.3% for influenza B.
Institutional influenza outbreaks ³	Similar	There were no new influenza outbreaks reported in Week 43, which was the same as the number of influenza outbreaks reported in Week 42 (0).
Influenza activity levels reported by health units	Lower	In Week 43, 1 public health unit reported 'localized' activity, 3 reported 'sporadic' activity, and 32 reported 'no activity'. In Week 42, 1 health unit reported 'localized' activity, 5 reported 'sporadic' activity, and 24 reported 'no activity'.
OVERALL ASSESSMENT	Similar	Overall, the indicators show that influenza activity for Week 43 was similar compared to Week 42.
Measures of severity for influenza		
Hospitalizations among laboratory-confirmed cases of influenza ⁴	There were 2 hospitalized cases reported in Week 43; the number of hospitalized cases reported in Week 42 was 5.	
Deaths among laboratory-confirmed cases of influenza ⁴	There were no deaths reported in Week 43 or in Week 42.	

Notes:

¹ Based on the date the case was reported to the public health unit.

² Positivity among specimens tested by Ontario laboratories that submit results to the Centre for Immunization and Respiratory Infectious Diseases (CIRID).

³ The number of new outbreaks reported for the current week is based on the date the outbreak was reported to the health unit; when reported date is unavailable, the date the outbreak was created in iPHIS is used

⁴ Hospitalization / death data are based on the date reported to the health unit. These data are likely under-reported and may not be comparable to prior years. Please see note on page 12.

Summary of respiratory pathogen activity for Surveillance Week 43, 2015

1. Circulating respiratory viruses in Ontario

Provincial percent positivity for both influenza A and B remains low, at 0.6% and 0.3% respectively in Week 43 ([Figure 1](#)). Rhinovirus was the most common circulating respiratory virus, with provincial positivity at 16.2% (Figures [2](#) and [3](#)).

2. Comparison to previous influenza seasons

For the 2015-2016 surveillance season to date, 76 laboratory-confirmed influenza cases have been reported. Among these cases, 85.5% (65/76) were influenza A ([Table 6](#)). Of the 36 reported influenza A cases with subtype information available, 80.6% (29/36) were H3N2 and 19.4% (7/36) were (H1N1)pdm09. Influenza activity in week 43 was low, which is within expected levels for this time of year.

3. Institutional respiratory outbreaks

No influenza outbreaks were reported in Week 43 ([Figure 6](#)). Influenza was detected in 1.1% (2/181) of institutional respiratory infection outbreaks reported to date in the 2015-2016 season. 59.7% (108/181) of the outbreaks were reported as entero/rhinovirus. Most respiratory infection outbreaks were reported from long-term care homes (105/181 or 58.0% of all reported outbreaks).

4. Severity - Influenza hospitalizations and deaths, and respiratory viruses by setting

In the season to date, 26 hospitalizations and no deaths have been reported among laboratory-confirmed influenza cases. The majority (24/26 or 92.3%) of hospitalizations occurred in influenza A cases, with the remaining hospitalizations (2/26 or 7.7%) in patients with combined influenza A and B. 57.7% (15/26) of the hospitalizations were in individuals 65 years of age and older. For week 43, influenza A was detected only from patients seen in community (ambulatory) settings or from institutional influenza outbreaks. Entero/rhinovirus was the respiratory virus most commonly detected from patients in community and institutional settings.

5. By Jurisdiction – Influenza activity

Influenza activity remains low across the province, with one public health unit reporting 'localized' activity, three health units reporting 'sporadic' activity, and 32 reporting 'no activity'.

6. By Age –Respiratory viruses

For week 43, influenza A was detected in one child in the 1-4 year age group and one individual in the 20-64 age group. Entero/rhinovirus was the most commonly detected virus among individuals 65 years and older. Information on influenza hospitalizations and deaths by age can be found in [Section 4: Severity - Influenza hospitalizations and deaths, and respiratory viruses by setting](#).

7. Strain comparisons – circulating influenza strains and vaccine strains

Through genetic characterization performed at the National Microbiology Laboratory (NML), sequence analysis showed that the one H1N1 virus tested nationally was antigenically similar to A/California/7/2009, which is the influenza A/H1N1 component [recommended by the World Health Organization](#) for the 2015-2016 Northern Hemisphere influenza vaccine.

Only one influenza A(H3N2) virus from Canada grew to sufficient titre for antigenic characterization by haemagglutination inhibition assay and was characterized as antigenically similar to the cell-passaged A/Switzerland/9715293/2013, the World Health Organization recommended influenza A(H3N2) component of the 2015-2016 Northern Hemisphere vaccine.

Of the six influenza B viruses characterized in Canada, NML reported that all were antigenically similar to the recommended influenza B component for the 2015-2016 vaccine, B/Phuket/3073/2013.

8. Antiviral resistance / sensitivity in influenza isolates

All influenza A viruses in Canada tested by the National Microbiology Laboratory (NML) for resistance in the 2015-2016 season were sensitive to oseltamivir and zanamivir, and resistant to amantadine. All influenza B viruses in Canada were sensitive to both oseltamivir and zanamivir.

9. Bacterial pathogens

Four specimens were positive for *Bordetella pertussis* and one specimen was positive for *Mycoplasma pneumoniae* during week 43. No specimen was positive for *Bordetella parapertussis*, *Bordetella holmesii* or *Chlamydomphila pneumoniae* during this week.


10. Data sources for this report

Provides a list and description of data sources used to prepare this report.

1. Circulating respiratory viruses in Ontario


Provincial percent positivity for both influenza A and B remains low, at 0.6% and 0.3% respectively in Week 43 ([Figure 1](#)). Rhinovirus was the most common circulating respiratory virus, with provincial positivity at 16.2% (Figures [2](#) and [3](#)).

Figure 1: Percentage of respiratory viral pathogens (influenza A, influenza B, respiratory syncytial virus, and parainfluenza virus) detected among specimens tested by all methods: Ontario, October 26, 2014 to October 31, 2015




Source: [Centre for Immunization and Respiratory Infectious Diseases \(CIRID\)](#)

Figure 2: Percentage of respiratory viral pathogens (adenovirus, human metapneumovirus, rhinovirus and coronavirus) detected among specimens tested by all methods: Ontario, October 26, 2014 to October 31, 2015



Source: [Centre for Immunization and Respiratory Infectious Diseases \(CIRID\)](#)

Figure 3: Total number of influenza tests performed and percent of positive tests by report week: Ontario, October 26, 2014 to October 31, 2015




Source: [Centre for Immunization and Respiratory Infectious Diseases \(CIRID\)](#)

2. Comparison to previous influenza seasons

For the 2015-2016 surveillance season to date, 76 laboratory-confirmed influenza cases have been reported. Among these cases, 85.5% (65/76) were influenza A ([Table 6](#)). Of the 36 reported influenza A cases with subtype information available, 80.6% (29/36) were H3N2 and 19.4% (7/36) were (H1N1)pdm09. Influenza activity in week 43 was low, which is within expected levels for this time of year.

Dominant Type/Subtype by Season	
2011-2012:	B (Yamagata lineage)
2012-2013:	A(H3N2)
2013-2014:	A(H1N1)pdm09
2014-2015:	A(H3N2)
2015-2016:	To Be Determined

Figure 4: Number of reported confirmed cases of influenza A by surveillance week: Ontario, September 1, 2011 to October 31, 2015




Source: [Integrated Public Health Information System \(iPHIS\)](#)

Notes:

- Interpret most recent case counts for the current season with caution due to reporting lags.
- Unlike the other seasons presented, the 2014-2015 season included a week 53; a week 53 occurs once every five to six years. Week 53 in 2014-15 corresponded to December 28, 2014 to January 3, 2015.

Figure 5: Number of reported confirmed cases of influenza B by surveillance week: Ontario, September 1, 2011 to October 31, 2015



Source: [Integrated Public Health Information System \(iPHIS\)](#)

Note:

- Interpret most recent case counts for the current season with caution due to reporting lags.
- Unlike the other seasons presented, the 2014-2015 season included a week 53; a week 53 occurs once every five to six years. Week 53 in 2014-15 corresponded to December 28, 2014 to January 3, 2015.

3. Institutional respiratory outbreaks

No influenza outbreaks were reported in Week 43 ([Figure 6](#)). Influenza was detected in 1.1% (2/181) of institutional respiratory infection outbreaks reported to date in the 2015-2016 season. 59.7% (108/181) of the outbreaks were reported as entero/rhinovirus. Most respiratory infection outbreaks were reported from long-term care homes (105/181 or 58.0% of all reported outbreaks).

Table 1: Institutional respiratory infection outbreaks: Ontario, Week 43 and total for the 2015-2016 season to date¹

Virus reported in outbreak	CURRENT WEEK		CUMULATIVE	
	Number of new outbreaks ²	Percentage of total for current week	Number of outbreaks cumulative	Percentage of total for cumulative
Influenza A ³	0	0.0%	2	1.1%
Influenza B ³	0	0.0%	0	0.0%
Both influenza A and B ³	0	0.0%	0	0.0%
Enterovirus/rhinovirus	1	7.1%	108	59.7%
Parainfluenza (all types)	1	7.1%	11	6.1%
Respiratory syncytial virus (RSV)	0	0.0%	0	0.0%
Human metapneumovirus, adenovirus, coronavirus or other viruses	0	0.0%	1	0.6%
Two or more non-influenza viruses ³	1	7.1%	9	5.0%
No organism reported	11	78.6%	50	27.6%
TOTAL	14	100.0%	181	100.0%

Source: [Integrated Public Health Information System \(iPHIS\)](#)

Notes:

¹ Season to date includes all outbreaks in which the date the outbreak was reported to the public health unit (or if unavailable, the date the outbreak was created in iPHIS) occurred between September 1, 2015 (Week 35, 2015) and the current reporting week, as well as outbreaks in which the reported date is missing but the outbreaks were entered into iPHIS during the current season.

² New outbreaks are defined as those in which the date the outbreak was reported to the health unit was in the current surveillance period, i.e. October 25, 2015 – October 31, 2015 (Week 43), and was recorded in iPHIS.

³ Any outbreak where influenza was identified is reported under the appropriate influenza category (“Influenza A”, “Influenza B”, or “Both influenza A and B”) regardless of what other virus was also identified in the outbreak.

Table 2: Institutional respiratory infection outbreaks by setting type: Ontario, total for the 2015-2016 season to date¹

Setting type reported	Number of influenza outbreaks (% of total)	Number of other respiratory viruses (% of total)
Long-Term Care Home	0 (0.0%)	105 (58.7%)
Hospital	0 (0.0%)	7 (3.9%)
Retirement Home	1 (50.0%)	29 (16.2%)
Other ²	0 (0.0%)	0 (0%)
Unknown	1 (50.0%)	38 (21.2%)
TOTAL	2 (100.0%)	179 (100.0%)


Source: [Integrated Public Health Information System \(iPHIS\)](#)

Notes:

¹ Season to date includes all outbreaks in which the date the outbreak was reported to the public health unit (or if unavailable, the date the outbreak was created in iPHIS) occurred between September 1, 2015 (Week 35, 2015) and the current reporting week, as well as outbreaks in which the reported date is missing but the outbreaks were entered into iPHIS during the current season.

² Other types of institutions include: correctional facilities, group homes, shelters, and facilities operating under the *Developmental Services Act*. Note that school-based and child care centre respiratory outbreaks are not captured in this table.

Figure 6: Institutional respiratory infection outbreaks by week of illness onset in the first case: Ontario, October 26, 2014 to October 31, 2015



Source: [Integrated Public Health Information System \(iPHIS\)](#)

Notes:

- Interpret most recent outbreak counts for the current season with caution due to reporting lags.
- Institutional respiratory infection outbreaks for which the date of onset of illness for the first case is missing are excluded in this figure. However, these outbreaks are counted in the cumulative outbreaks section of Table 1.
- Any outbreak where influenza was identified is reported under the appropriate influenza category ("Influenza A", "Influenza B", or "Both influenza A & B") regardless of what other virus is also identified in the outbreak.

4. Severity - Influenza hospitalizations and deaths, and respiratory viruses by setting

In the season to date, 26 hospitalizations and no deaths have been reported among laboratory-confirmed influenza cases. The majority (24/26 or 92.3%) of hospitalizations occurred in influenza A cases, with the remaining hospitalizations (2/26 or 7.7%) in patients with combined influenza A and B. 57.7% (15/26) of the hospitalizations were in individuals 65 years of age and older.

For week 43, influenza A was detected only from patients seen in community (ambulatory) settings or from institutional influenza outbreaks. Enterovirus was the respiratory virus most commonly detected from patients in community and institutional settings.

Table 3: Hospitalizations and deaths among confirmed influenza cases by age group: Ontario, Week 43 and cumulative for the 2015-2016 influenza season

Age Group	HOSPITALIZATIONS		DEATHS	
	Weekly Count	Cumulative count (rate per 100,000)	Weekly count	Cumulative count (rate per 100,000)
<1	0	0 (0.0)	0	0 (0.0)
1 – 4	1	1 (0.2)	0	0 (0.0)
5 – 14	0	0 (0.0)	0	0 (0.0)
15 – 24	0	1 (0.1)	0	0 (0.0)
25 – 44	0	3 (0.1)	0	0 (0.0)
45 – 64	1	6 (0.2)	0	0 (0.0)
65+	0	15 (0.7)	0	0 (0.0)
Total	2	26 (0.2)	0	0 (0.0)

Source: [Integrated Public Health Information System \(iPHIS\)](#) and [IntelliHEALTH Ontario](#)

Notes:

- Reported deaths in any laboratory-confirmed influenza case are included; the contribution of influenza to the death is not able to be determined.
- The numbers of hospitalizations and deaths provided in iPHIS need to be interpreted with caution due to delays between the onset of illness, hospital admission or death, and when this information is reported to public health. During periods of increased influenza activity, there may be additional delays in the reporting of hospitalizations and deaths beyond the expected reporting time lag.
- The numbers of hospitalizations and deaths reported in iPHIS are likely an underestimation of the true experience in the Ontario population for many reasons, including underreporting because influenza is not suspected as the source of infection so testing is not done.
- In addition, public health units were advised in December 2014 and August 2015 that only 20% of laboratory confirmed cases require follow-up, as opposed to all cases. As a result, there will likely be more under-reporting of hospitalization and deaths in the 2014-2015 and 2015-2016 seasons than in previous seasons so comparisons to prior years should be done with caution.

Table 4: Number and percentage of specimens (in brackets)¹ with respiratory pathogens detected by patient setting, week 43 and cumulative for the 2015-2016 influenza season

Respiratory Pathogen Detected	Patient setting when tested									
	Intensive Care Unit (ICU) Number (% positive) ¹		Hospital (Non-ICU) Number (% positive) ¹		Emergency Number (% positive) ¹		Ambulatory/No setting Number (% positive) ¹		Institution ² Number (% positive) ¹	
	Current	Cumulative	Current	Cumulative	Current	Cumulative	Current	Cumulative	Current	Cumulative
Influenza A (all) ³	0 (0.0)	0 (0.0)	0 (0.0)	4 (0.4)	0 (0.0)	39 (4.4)	2 (1.2)	18 (1.2)	0 (0.0)	3 (0.7)
Influenza A/ (H3) ⁴	0	0	0	3	0	23	0	10	0	2
Influenza A/(H1N1)pdm09 ⁴	0	0	0	1	0	11	0	1	0	0
Influenza B	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.1)	0 (0.0)	4 (0.3)	0 (0.0)	0 (0.0)
Adenovirus	0 (0.0)	2 (0.7)	1 (0.7)	6 (0.6)	3 (2.5)	18 (2.0)	5 (3.1)	24 (1.6)	0 (0.0)	0 (0.0)
Coronavirus	0 (0.0)	1 (0.4)	0 (0.0)	0 (0.0)	0 (0.0)	8 (10.8)	0 (0.0)	2 (0.5)	0 (0.0)	1 (0.3)
Enterorhinovirus	3 (16.7)	54 (19.6)	3 (42.9)	23 (16.7)	0 (0.0)	22 (17.3)	5 (25.0)	112 (22.4)	10 (71.4)	190 (63.3)
Human metapneumovirus	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.1)	0 (0.0)	2 (0.2)	1 (0.6)	5 (0.4)	0 (0.0)	1 (0.2)
Parainfluenza(all types)	1 (4.8)	8 (2.8)	6 (4.3)	32 (3.2)	4 (3.4)	37 (4.2)	7 (4.3)	55 (3.7)	1 (4.5)	18 (4.1)
Respiratory syncytial virus	0 (0.0)	0 (0.0)	2 (1.4)	8 (0.8)	0 (0.0)	1 (0.1)	1 (0.6)	5 (0.3)	1 (4.5)	2 (0.5)
Total positives⁵	4 (16.7)	65 (22.3)	12 (8.4)	74 (7.3)	7 (5.8)	128 (14.5)	21 (12.7)	225 (15.3)	12 (54.5)	215 (48.8)

Source: [Public Health Ontario Laboratory \(PHOL\)](#)

Notes:

¹ The percentages in this table are calculated as number of specimens with the respiratory pathogen detected divided by total number of specimens tested for that particular pathogen within the patient setting. For example, if 40 respiratory specimens were tested for influenza B from patients in intensive care unit (ICU) settings and 4 tested positive, the cell for influenza B in the ICU setting would indicate “4 (10%)”.

² Institution includes retirement homes, correctional facilities, and undefined institutions

³ The “Influenza A (all)” row includes the numbers in the “Influenza A (H3)” and “Influenza A/(H1N1)pdm09” rows as well as Influenza A positives that were not subtyped.

⁴ Percent positivity is not calculated for Influenza A (H3) and Influenza A/(H1N1)pdm09 because subtyping is only done on select influenza A samples, precluding an accurate calculation.

⁵ When calculating the “Total Positives”, the “Influenza A (H3)” and “Influenza A/(H1N1)pdm09” rows are not counted as these numbers are already included in the “Influenza A (all)” row.

Notes:

- **Table 5** is information from Public Health Ontario Laboratories (PHOL). It allows for the calculation of percent positivity because PHOL tests the samples and knows the numbers of positive and negative results. Percent positivity for health units that had less than 40 respiratory specimens tested may not be reliable due to small numbers.
- **Table 6** is information from the Integrated Public Health Information System (iPHIS) based on positive influenza cases reported to the health unit. These tests are done by multiple laboratories and because negative results are not reported to the health unit, the percent positivity cannot be calculated.

Legend

Influenza Level

- No Data
- No Activity
- Sporadic
- Localized
- Widespread

Health Regions and Units

Central East	Eastern	South West
DUR Durham Region	EOH Eastern Ontario	CHK Chatham-Kent
HKP Haliburton, Kawartha, Pine Ridge	HPE Hastings-Prince Edward	ELG Elgin-St. Thomas
PEE Peel Region	KFL Kingston-Frontenac	GBO Grey-Bruce
PTC Peterborough County-City	LGL Leeds-Grenville-Lanark	HUR Huron County
SMD Simcoe-Muskoka	OTT City of Ottawa	LAM Lambton
YRK York Region	REN Renfrew County	MSL Middlesex-London
Central West	North East	OXF Oxford County
BRN Brant	ALG Algoma	PDH Perth County
HDN Haldimand-Norfolk	NPS North Bay - Parry Sound	WEC Windsor-Essex
HAL Halton Region	PQP Porcupine	Toronto
HAM City of Hamilton	SUD Sudbury and District	TOR Toronto
NIA Niagara Region	TSK Timiskaming	
WAT Waterloo Region	North West	
WDG Wellington-Dufferin-Guelph	NWR Northwestern	
	THB Thunder Bay	

Page 14 of 23

Table 5: Specimens submitted to Public Health Ontario Laboratories for influenza testing and positive results by public health unit; Number for week 43 and cumulative total for the 2015-2016 influenza season (in brackets)

Health Unit and Region	Number of specimens submitted	Number of specimen tested	Influenza A				Influenza B	
			Number of influenza A positive specimens	H3	A(H1N1) pdm09	% positivity influenza A	Number of influenza B positive specimens	% positivity influenza B
Northwestern	8 (74)	4 (70)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Thunder Bay District	5 (66)	5 (62)	0 (1)	0 (0)	0 (0)	0 (2)	0 (0)	0 (0)
TOTAL NORTH WEST	13 (140)	9 (132)	0 (1)	0 (0)	0 (0)	0 (1)	0 (0)	0 (0)
Algoma	4 (40)	3 (39)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
North Bay Parry Sound District	16 (71)	12 (66)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Porcupine	8 (51)	7 (49)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Sudbury & District	11 (51)	10 (49)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Timiskaming	2 (25)	2 (25)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
TOTAL NORTH EAST	41 (238)	34 (228)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
City of Ottawa	1 (55)	0 (49)	0 (1)	0 (0)	0 (0)	NA (0)	0 (0)	NA (0)
Eastern Ontario	11 (57)	7 (53)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Hastings & Prince Edward Counties	4 (54)	4 (53)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Kingston, Frontenac, Lennox & Addington	22 (81)	13 (72)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Leeds, Grenville And Lanark District	1 (29)	1 (29)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Renfrew County And District	2 (17)	1 (16)	0 (0)	0 (0)	0 (0)	0 (0)	0 (1)	0 (6)
TOTAL EASTERN	41 (293)	26 (272)	0 (1)	0 (0)	0 (0)	0 (0)	0 (1)	0 (0)
Durham Region	37 (286)	27 (264)	0 (10)	0 (9)	0 (0)	0 (4)	0 (0)	0 (0)
Haliburton, Kawartha, Pine Ridge	20 (110)	13 (103)	0 (2)	0 (2)	0 (0)	0 (2)	0 (0)	0 (0)
Peel Region	68 (483)	57* (466)	1 (19)	0 (7)	0 (8)	2 (4)	0 (3)	0 (1)
Peterborough County-City	7 (56)	6 (53)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Simcoe Muskoka District	61 (367)	28 (328)	0 (6)	0 (5)	0 (0)	0 (2)	0 (0)	0 (0)
York Region	43 (262)	28 (239)	0 (5)	0 (2)	0 (3)	0 (2)	0 (0)	0 (0)
TOTAL CENTRAL EAST	236 (1564)	159 (1453)	1 (42)	0 (25)	0 (11)	1 (0)	0 (3)	0 (0)
Toronto	150 (971)	115* (907)	0 (8)	0 (6)	0 (1)	0 (1)	0 (0)	0 (0)
TOTAL TORONTO	150 (971)	115 (907)	0 (8)	0 (6)	0 (1)	0 (1)	0 (0)	0 (0)
Chatham-Kent	6 (27)	3 (23)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Elgin-St. Thomas	6 (38)	3 (34)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Grey Bruce	8 (65)	7 (62)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Huron County	1 (16)	1 (16)	0 (1)	0 (1)	0 (0)	0 (6)	0 (0)	0 (0)
Lambton County	6 (43)	4 (41)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Middlesex-London	5 (49)	5 (49)	0 (2)	0 (2)	0 (0)	0 (4)	0 (0)	0 (0)
Oxford County	3 (30)	3 (29)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Perth District	4 (31)	4 (29)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)

Windsor-Essex County	17 (134)	13 (129)	0 (1)	0 (1)	0 (0)	0 (1)	0 (0)	0 (0)
TOTAL SOUTH WEST	56 (433)	43 (412)	0 (4)	0 (4)	0 (0)	0 (1)	0 (0)	0 (0)
Brant County	9 (95)	7 (93)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
City Of Hamilton	6 (62)	3 (56)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Haldimand-Norfolk	13 (33)	9 (29)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Halton Region	28 (132)	22 (121)	0 (2)	0 (0)	0 (1)	0 (2)	0 (0)	0 (0)
Niagara Region	29 (138)	21 (127)	0 (2)	0 (2)	0 (0)	0 (2)	0 (1)	0 (1)
Waterloo Region	19 (136)	14 (130)	1 (4)	0 (1)	0 (0)	7 (3)	0 (0)	0 (0)
Wellington-Dufferin-Guelph	11 (113)	10 (110)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
TOTAL CENTRAL WEST	115 (709)	86 (666)	1 (8)	0 (3)	0 (1)	1 (1)	0 (1)	0 (0)
OUT OF PROVINCE	4 (35)	3 (34)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
TOTAL ONTARIO	656 (4383)	475 (4104)	2 (64)	0 (38)	0 (13)	0 (2)	0 (5)	0 (0)

Notes:

- Cumulative numbers are from September 1, 2015 to the current week (week 43, 2015) and are displayed in brackets.
- The difference between submissions and testing is due to lags in testing time.
- "Health Unit" reflects the jurisdiction where the patient resides. In the event this is not available, the health unit of the specimen submitter is used.
- *Indicates health units that tested more than 40 specimens tested during the current week. Percent positivity for health units that had less than 40 respiratory specimens tested may not be reliable due to small numbers.
- NA-Not applicable- Indicates Health units with no specimens tested; therefore percent positivity was not calculated.

Source: [Public Health Ontario Laboratory \(PHOL\)](#)

Table 6. Number of reported confirmed influenza cases by public health unit and health region; Number for week 43 and cumulative total for the 2015-2016 influenza season (in brackets)

Health Unit and Region	Number of influenza A positive cases			Influenza A & B	Influenza B	TOTAL
	(H1N1) pdm09	H3	Influenza A All subtypes ¹			
Northwestern	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Thunder Bay District	0 (0)	0 (0)	0 (1)	0 (0)	0 (0)	0 (1)
TOTAL NORTH WEST	0 (0)	0 (0)	0 (1)	0 (0)	0 (0)	0 (1)
Algoma	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
North Bay Parry Sound District	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Porcupine	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Sudbury & District	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Timiskaming	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
TOTAL NORTH EAST	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
City of Ottawa	0 (0)	0 (0)	0 (5)	0 (0)	0 (0)	0 (5)
Eastern Ontario	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Hastings & Prince Edward Counties	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Kingston, Frontenac, Lennox & Addington	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Leeds, Grenville And Lanark District	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Renfrew County And District	0 (0)	0 (0)	0 (0)	0 (0)	0 (1)	0 (1)
TOTAL EASTERN	0 (0)	0 (0)	0 (5)	0 (0)	0 (1)	0 (6)
Durham Region	0 (0)	0 (6)	0 (6)	0 (0)	0 (0)	0 (6)
Haliburton, Kawartha, Pine Ridge	0 (0)	0 (2)	0 (2)	0 (0)	0 (0)	0 (2)
Peel Region	0 (3)	1 (4)	2 (15)	0 (0)	4 (6)	6 (21)
Peterborough County-City	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Simcoe Muskoka District	0 (0)	0 (6)	1 (7)	0 (0)	0 (0)	1 (7)
York Region	0 (1)	0 (2)	0 (5)	0 (0)	0 (0)	0 (5)
TOTAL CENTRAL EAST	0 (4)	1 (20)	3 (35)	0 (0)	4 (6)	7 (41)
Toronto	0 (2)	1 (5)	2 (8)	0 (1)	0 (2)	2 (11)
TOTAL TORONTO	0 (2)	1 (5)	2 (8)	0 (1)	0 (2)	2 (11)
Chatham-Kent	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Elgin-St. Thomas	0 (0)	0 (0)	0 (1)	0 (0)	0 (0)	0 (1)
Grey Bruce	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Huron County	0 (0)	0 (0)	0 (1)	0 (0)	0 (0)	0 (1)
Lambton County	0 (0)	0 (0)	0 (1)	0 (0)	0 (0)	0 (1)
Middlesex-London	0 (0)	0 (2)	0 (4)	0 (0)	0 (0)	0 (4)
Oxford County	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Perth District	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Windsor-Essex County	0 (0)	0 (1)	0 (1)	0 (0)	0 (0)	0 (1)
TOTAL SOUTH WEST	0 (0)	0 (3)	0 (8)	0 (0)	0 (0)	0 (8)
Brant County	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
City Of Hamilton	0 (0)	0 (0)	0 (1)	0 (0)	0 (0)	0 (1)
Haldimand-Norfolk	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Halton Region	0 (1)	0 (0)	0 (3)	0 (0)	0 (0)	0 (3)
Niagara Region	0 (0)	0 (1)	0 (1)	0 (0)	0 (1)	0 (2)
Waterloo Region	0 (0)	0 (0)	0 (3)	0 (0)	0 (0)	0 (3)
Wellington-Dufferin-Guelph	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
TOTAL CENTRAL WEST	0 (1)	0 (1)	0 (8)	0 (0)	0 (1)	0 (9)
TOTAL ONTARIO	0 (7)	2 (29)	5 (65)	0 (1)	4 (10)	9 (76)

Source: [Integrated Public Health Information System \(iPHIS\)](#)

Note:

¹ Includes influenza A positive cases for whom no subtype was reported

[BACK TO TOP](#)

6. By Age –Respiratory viruses

For week 43, influenza A was detected in one child in the 1-4 year age group and one individual in the 20-64 age group. Enterovirus was the most commonly detected virus among individuals 65 years and older. Information on influenza hospitalizations and deaths by age can be found in [Section 4: Severity - Influenza hospitalizations and deaths, and respiratory viruses by setting](#).

Table 7: Number and percentage of specimens (in brackets)¹ with respiratory pathogens detected by age group by Public Health Ontario Laboratories, week 43 and cumulative for the 2015-2016 influenza season

Respiratory Pathogen Detected	Age groups									
	<1 Number (% positive) ¹		1-4 Number (% positive) ¹		5-19 Number (% positive) ¹		20-64 Number (% positive) ¹		65+ Number (% positive) ¹	
	Current	Cumulative	Current	Cumulative	Current	Cumulative	Current	Cumulative	Current	Cumulative
Influenza A (all) ²	0 (0.0)	3 (0.5)	1 (5.9)	1 (0.5)	0 (0.0)	3 (1.5)	1 (0.6)	34 (2.4)	0 (0.0)	23 (1.3)
Influenza A (H3) ³	0	2	0	0	0	1	0	18	0	17
Influenza A(H1N1)pdm09 ³	0	1	0	0	0	1	0	9	0	2
Influenza B	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	4 (0.3)	0 (0.0)	1 (0.1)
Adenovirus	1 (1.6)	15 (2.7)	2 (11.8)	11 (5.9)	2 (6.5)	6 (3.0)	4 (2.3)	15 (1.1)	0 (0.0)	3 (0.2)
Coronavirus	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	5 (1.3)	0 (0.0)	7 (1.1)
Enterovirus	1 (50.0)	9 (14.3)	0 (0.0)	1 (9.1)	2 (66.7)	6 (19.4)	2 (12.5)	84 (16.7)	16 (42.1)	301 (41.1)
Human metapneumovirus	0 (0.0)	3 (0.5)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.6)	3 (0.2)	0 (0.0)	3 (0.2)
Parainfluenza	8 (12.5)	53 (9.4)	2 (11.8)	22 (11.8)	1 (3.2)	4 (2.0)	2 (1.2)	19 (1.4)	6 (3.4)	52 (3.0)
Respiratory syncytial virus	1 (1.6)	9 (1.6)	1 (5.9)	1 (0.5)	0 (0.0)	0 (0.0)	1 (0.6)	1 (0.1)	1 (0.6)	5 (0.3)
Total positives⁴	11 (16.7)	92 (16.2)	6 (35.3)	36 (19.3)	5 (15.6)	19 (9.3)	11 (6.1)	165 (11.9)	23 (12.7)	395 (22.5)

Source: [Public Health Ontario Laboratory \(PHOL\)](#)

Notes:

¹ The percent positivity in this table is calculated as number of specimens with the respiratory pathogen detected divided by total number of specimens tested for that particular pathogen within the age group. For example, if 50 respiratory specimens were tested for influenza B from patients in the 5-19 year age group and 10 tested positive, the cell for influenza B in 5-19 years olds would indicate "10 (20%)".

² The "Influenza A (all)" row includes the numbers in the "Influenza A (H3)" and "Influenza A/(H1N1)pdm09" rows as well as Influenza A positives that were not subtyped.

³ Percent positivity is not calculated for Influenza A (H3) and "Influenza A/(H1N1)pdm09" because subtyping is only done on select influenza A samples, precluding an accurate calculation.

⁴ When calculating the "Total Positives", the "Influenza A (H3)" and "Influenza A/(H1N1)pdm09" rows are not counted as these numbers are already included in the "Influenza A (all)" row.

[BACK TO TOP](#)

7. Strain comparisons – circulating influenza strains and vaccine strains

Through genetic characterization performed at the National Microbiology Laboratory (NML), sequence analysis showed that the one H1N1 virus tested nationally was antigenically similar to A/California/7/2009, which is the influenza A/H1N1 component [recommended by the World Health Organization](#) for the 2015-2016 Northern Hemisphere influenza vaccine.

Only one influenza A(H3N2) virus from Canada grew to sufficient titre for antigenic characterization by haemagglutination inhibition assay and was characterized as antigenically similar to the cell-passaged A/Switzerland/9715293/2013, the World Health Organization recommended influenza A(H3N2) component of the 2015-2016 Northern Hemisphere vaccine.

Of the six influenza B viruses characterized in Canada, NML reported that all were antigenically similar to the recommended influenza B component for the 2015-2016 vaccine, B/Phuket/3073/2013.

Table 8: Strain characterization completed on influenza isolates at the National Microbiology Laboratory: Ontario and Canada, September 1, 2015 to November 5, 2015¹

Influenza strains	Ontario	Canada
Influenza A (H3N2) A/Switzerland/9715293/13-like	1	1
Influenza A (H1N1) A/California/07/09-like	0	1
Influenza B B/Phuket/3073/13-like	1	6

Source: [National Microbiology Laboratory \(NML\)](#)

[BACK TO TOP](#)

8. Antiviral resistance / sensitivity for influenza

All influenza A viruses in Canada tested by the National Microbiology Laboratory (NML) for resistance in the 2015-2016 season were sensitive to oseltamivir and zanamivir, and resistant to amantadine. All influenza B viruses in Canada were sensitive to both oseltamivir and zanamivir.

Table 9: Amantadine, oseltamivir and zanamivir susceptibility assays completed on influenza isolates at the National Microbiology Laboratory: Ontario and Canada, September 1, 2015 to November 5, 2015

Influenza strains	Amantadine				Oseltamivir				Zanamivir			
	ONTARIO		CANADA		ONTARIO		CANADA		ONTARIO		CANADA	
	R	S	R	S	R	S	R	S	R	S	R	S
Influenza A (H3N2)	6	0	21	0	0	4	0	18	0	4	0	18
Influenza A (H1N1)pdm09	0	0	1	0	0	0	0	1	0	0	0	1
Influenza B	NA	NA	NA	NA	0	1	0	5	0	1	0	5

Source: [National Microbiology Laboratory \(NML\)](#)

R = Resistant, S = Susceptible, NA = Not Applicable

[BACK TO TOP](#)

9. Bacterial pathogens

Four specimens were positive for *Bordetella pertussis* and one specimen was positive for *Mycoplasma pneumoniae* during week 43. No specimen was positive for *Bordetella parapertussis*, *Bordetella holmesii* or *Chlamydophila pneumoniae* during this week.

Table 10: Number and percent positivity of respiratory specimens tested for any *Bordetella* species, *Mycoplasma pneumoniae* or *Chlamydophila pneumoniae*, Public Health Ontario Laboratories, week 43

Pathogens	Tested	Positives	% positivity
	Current (Cumulative)	Current (Cumulative)	Current (Cumulative)
<i>Bordetella pertussis</i>	80 (3250)	4 (238)	5 (7)
<i>Bordetella parapertussis</i>	80 (3250)	0 (14)	0 (0)
<i>Bordetella holmesii</i>	80 (3250)	0 (3)	0 (0)
<i>Mycoplasma pneumoniae</i>	11 (417)	1 (22)	9 (5)
<i>Chlamydophila pneumoniae</i>	11 (417)	0 (2)	0 (1)

Source: [Public Health Ontario Laboratory \(PHOL\)](#)

Notes:

- Cumulative numbers (in brackets) for *Bordetella* are from January 1, 2015.
- Cumulative numbers (in brackets) for *Mycoplasma* and *Chlamydophila* are from September 1, 2015.

10. Data sources for this report

- **Integrated Public Health Information System (iPHIS):** provides information on laboratory-confirmed cases of influenza reported to local public health units, and institutional outbreaks of influenza and other respiratory pathogens reported to local public health units. iPHIS is administered by the Ontario Ministry of Health and Long-Term Care; data for this issue was extracted by Public Health Ontario on November 4, 2015.

Case counts for influenza A and B cases are based on the 'Reported Date', which was the date the health unit was notified of the case; however, cases are assigned to a particular surveillance week based on the episode date entered in iPHIS for the case. Episode date for a case corresponds to the earliest date on record for the case according to the iPHIS hierarchy (Symptom Date > Clinical Diagnosis Date > Specimen Collection Date > Lab Test Date > Reported Date). Cases are excluded from these counts if the episode date is based on reported date, and the reported date occurs after the latest surveillance week.

- **Centre for Immunization and Respiratory Infectious Diseases (CIRID) of the Public Health Agency of Canada (PHAC):** provides information on the number of influenza tests and number that are positive based on submissions from 16 participating laboratories in Ontario including 11 Public Health Ontario Laboratories (PHOLs) and five hospital-based laboratories. The results are assigned to a particular surveillance week based on when test results are reported to PHAC; these data are not updated when results are submitted late for previous surveillance weeks. These data represent the number of specimens tested, which may not necessarily correspond with the number of patients as more than one specimen may have been submitted per patient. Cumulative numbers for the season to date are also available through [FluWatch](#).
- **Public Health Ontario Laboratory (PHOL):** provides specimen level information from the 11 Public Health Ontario Laboratory sites located in Ontario; data for the issue was extracted from the Laboratory Information Management System by Public Health Ontario on November 4, 2015
- **National Microbiology Laboratory (NML), Influenza and Respiratory Viruses Section:** provides testing results for influenza strain identification and influenza antiviral resistance / susceptibility
- **Public Health Ontario from public health units:** Public health units provide Public Health Ontario with weekly reports of influenza activity levels in their areas [Provincial Influenza Activity Report (Appendix C) Database]. Influenza activity levels are assigned by local public health units and reported to Public Health Ontario by 4:00 pm on the Tuesday following the end of each surveillance week. Activity levels are assigned based on laboratory confirmations, ILI reports from various sources, and laboratory-confirmed institutional respiratory infection outbreaks. Please refer to the [detailed definitions for the 2015-2016 season](#) for more information. Activity levels reported for a particular surveillance week may not necessarily correspond to the number of new outbreaks reported in the same week in Table 3 because ongoing outbreaks from previous weeks, as well as laboratory confirmed outbreaks in schools, may be included in the assessment of the activity level.

- **IntelliHEALTH Ontario:** provides information on population counts based on 2013 estimates. The information was obtained from the Ontario Ministry of Health and Long-Term Care, and extracted by Public Health Ontario on July 3, 2014.