
Frost Injury in Sorghum Species

Posted on [September 27, 2023](#) by [Christine O'Reilly, OMAFRA Forage & Grazing Specialist](#)

Caution! Members of the sorghum family – sorghum, sudangrass, and hybrid sorghum-sudangrass – contain dhurrin, a glucoside that breaks down to release hydrocyanic acid, also known as prussic acid (hydrogen cyanide, HCN). A sudden disruption of growth such as frost, drought or cutting, causes prussic acid to be released inside the plant at a more rapid rate. High prussic acid levels may be lethal to ruminants; death often occurs within 20 minutes of ingesting sorghum forage with elevated prussic acid concentrations.

Early fall, **between the first frost and the first killing frost, is the riskiest time of year for prussic acid** production in sorghum species. These warm-season grasses are very frost-sensitive. It is impossible to know by looking at them how much prussic acid is present in the plants. Below are some frequently asked questions related to managing sorghum species and frost.

Is it okay to harvest sorghum after a frost?

Wait 3-5 days after a killing frost to ensile sorghum or make baleage. Prussic acid levels will decrease with fermentation. Silage should be completely fermented before feeding (6–8 weeks).

I cut my sorghum, then we got a frost while it was in the swath. Will the sorghum still produce prussic acid?

Cut grasses don't know they have been cut. Cellular processes, like respiration and photosynthesis, continue so long as the leaves have enough water and oxygen to keep going. While information on this specific situation is scarce, the safest option is to respect the same wait times as if the crop was hit by frost just before cutting. This will give the prussic acid time to dissipate.

There is frost in the forecast, and my cattle/sheep are grazing sorghum. Should I move them off the sorghum?

Yes. Prussic acid levels are highest just after the frost occurs. Since this is usually in the wee hours of the morning, it is generally much easier to avoid an issue by removing livestock from the sorghum before the forecasted frost.

Sometimes frost comes as a surprise. Immediately after a frost, remove the livestock from the pasture until it has dried out (usually 6–7 days). If new shoots develop, harvest the field as silage rather than pasture.

Prussic acid concentration is always higher in leaves than in stems. Because livestock preferentially graze leaves, they are more likely to be poisoned by prussic acid in a pasture situation. Lightly grazing a field will not reduce risk.

References

Cassida, K. 2019. Frosted sorghum and sudangrass pasture poses prussic acid poisoning risk. Michigan State University. Retrieved 21 September 2020 from https://www.canr.msu.edu/news/frosted_sorghum-sudangrass_pasture_poses_prussic_acid_poisoning_risk

Vough, L. Date Unknown. Preventing prussic acid poisoning of livestock. Oregon State University. Retrieved 21 September 2020 from <https://forages.oregonstate.edu/fi/topics/pasturesandgrazing/grazingsystemdesign/preventingprussicacidpoisoning>

Sponsored Content



Taking Viagra? Use This Household Food To Fix ED

urolotrck.com | Sponsored

Intimate Clothing Trends in 2023 May Surprise You

Bras | Sponsored

Ontario Pensioners Born Before 1967 Are Eligible For Invisible Hearing Aids

Best Hearing Aids | Sponsored

Urologist: 87% of Men with E.D. Don't Know About This Easy Solution (Try Tonight!)

urologytip.pro | Sponsored

Seniors In Ontario Eligible For Hearing Aids In Fall 2023

Connect Hearing | Sponsored

[Gallery] It's No Secret Why We Don't See This Actress Anymore

Investing Magazine | Sponsored

Toronto: Claim This Benefit Before The End of 2023


canadianpayout.com | Sponsored

Ontario Seniors Born Before 1967 Are Eligible For New Hearing Aids

www.besthearingaids.ca | Sponsored

Toronto Residents Should Claim This Benefit

financialhackz.com | Sponsored

**About Christine O'Reilly, OMAFRA Forage & Grazing Specialist**

Areas of Focus: - Production and storage strategies to improve the productivity and profitability of forages - Benefits of forages and grazing within cropping systems - Forage production and grazing systems in northern Ontario

[View all posts by Christine O'Reilly, OMAFRA Forage & Grazing Specialist →](#)

This entry was posted in [Crops](#), [Livestock](#), [Weather](#) and tagged [Forage](#). Bookmark the [permalink](#).

ONOrganic

Create a free website or blog at WordPress.com.