

EPIDEMIOLOGICAL SUMMARY

MPOX in Ontario: May 1, 2022 to November 29, 2022

Introduction

This report provides an epidemiologic summary of MPOX (formerly known as monkeypox) in Ontario and includes the most current information available from Ontario's integrated Public Health Information System (iPHIS) as of **November 29, 2022**.

The current provincial case definition for MPOX can be found in Appendix 1 of the Infectious Disease Protocol for [Smallpox and other Orthopoxviruses including Monkeypox](#).


For further information regarding the MPOX virus, including signs, symptoms, and how to reduce your risk of infection, visit the Ontario Ministry of Health's [Monkeypox Virus](#) webpage and Public Health Ontario's [MPOX webpage](#) for additional resources.

Highlights

- There are 691 confirmed cases of MPOX in Ontario (1 new confirmed case since the last report).
 - 505/691 (73.1%) confirmed cases were reported by Toronto Public Health (see [Table 3](#) for confirmed cases by public health unit).
 - 685/691 (99.1%) confirmed cases are male and 6/691 (0.9%) are female.
 - The average age of confirmed cases is 36.6 years (range: <20 – 74 years).
 - The most commonly reported symptoms include rash, oral/genital lesions, swollen lymph nodes, headache, fever, chills, myalgia and fatigue.
 - 20/691 (2.9%) of confirmed cases have been hospitalized, and 2/691 (0.3%) have been in the intensive care unit (ICU); no deaths have been reported.
- In addition, there are 10 probable cases. 10/10 (100.0%) are male; the 10 probable cases range in age from 21 to 69 years (average: 43.2 years).


Case Characteristics

Figure 1. MPOX cases by accurate episode date and case classification



Note: Accurate Episode Date is defined in the [Data Caveats](#). Due to potential delays between symptom onset, reporting of laboratory test results, and data entry into iPHIS, case counts for the last 10 days (gray area) may be incomplete and should be interpreted with caution.

Figure 2. Number of individuals tested for MPOX and percent positivity by week



Data Source: Public Health Ontario Laboratory Information Management System.

Note: Week was assigned using sample collection date, if provided, and login date otherwise. Not all testing has been completed for the most recent week.

Table 1. Gender* and age group (in years) of MPOX cases by case classification

Gender and Age Group	Confirmed n (%)	Probable n (%)	Total n (%)
Gender: Male	685 (99.1)	10 (100.0)	695 (99.1)
Gender: Female	6 (0.9)	0 (0.0)	6 (0.9)
Gender: Unknown	0 (0.0)	0 (0.0)	0 (0.0)
Age group: < 20 years	5 (0.7)	0 (0.0)	5 (0.7)
Age group: 20 - 29 years	158 (22.9)	1 (10.0)	159 (22.7)
Age group: 30 - 39 years	289 (41.8)	4 (40.0)	293 (41.8)
Age group: 40 - 49 years	168 (24.3)	3 (30.0)	171 (24.4)
Age group: 50 - 59 years	53 (7.7)	0 (0.0)	53 (7.6)
Age group: 60 - 69 years	14 (2.0)	2 (20.0)	16 (2.3)
Age group: ≥ 70 years	4 (0.6)	0 (0.0)	4 (0.6)
Age group: Unknown	0 (0.0)	0 (0.0)	0 (0.0)
Total	691 (98.6)	10 (1.4)	701 (100.0)

***Note:** The gender category captures an individual's internal and individual experience of gender and not necessarily their sex assignment at birth. Case counts may fluctuate based on data cleaning at the local public health unit level.

Table 2. Predominant symptoms reported by MPOX cases by case classification

Symptom	Confirmed n (%)	Probable n (%)	Total n (%)
Rash	532 (77.0)	9 (90.0)	541 (77.2)
Oral/genital lesions	425 (61.6)	8 (80.0)	433 (61.8)
Fever	332 (48.0)	5 (50.0)	337 (48.1)
Fatigue	305 (44.1)	5 (50.0)	310 (44.2)
Swollen lymph nodes	300 (43.4)	3 (30.0)	303 (43.2)
Chills	288 (41.7)	5 (50.0)	293 (41.8)
Headache	223 (32.3)	3 (30.0)	226 (32.2)
Myalgia	217 (31.4)	3 (30.0)	220 (31.4)
Sweating	184 (26.6)	3 (30.0)	187 (26.7)
Sore throat	166 (24.0)	2 (20.0)	168 (24.0)
Prostration (exhaustion)	120 (17.4)	2 (20.0)	122 (17.4)
Back pain/back ache	103 (14.9)	3 (30.0)	106 (15.1)
Other*	235 (34.0)	5 (50.0)	240 (34.2)

***Note:** other symptoms include: constipation, coryza (runny nose), cough, diarrhea, dysuria (painful urination), malaise, and nausea.

Table 3. MPOX cases by diagnosing public health unit and case classification

Public Health Unit	Confirmed n (%)	Probable n (%)	Total n (%)
Algoma Public Health	0 (0.0)	0 (0.0)	0 (0.0)
Brant County Health Unit	4 (0.6)	0 (0.0)	4 (0.6)
Chatham-Kent Public Health	0 (0.0)	0 (0.0)	0 (0.0)
City of Hamilton Public Health Services	20 (2.9)	0 (0.0)	20 (2.9)
Durham Region Health Department	11 (1.6)	0 (0.0)	11 (1.6)
Eastern Ontario Health Unit	0 (0.0)	0 (0.0)	0 (0.0)
Grey Bruce Health Unit	2 (0.3)	1 (10.0)	3 (0.4)
Haldimand-Norfolk Health Unit	3 (0.4)	0 (0.0)	3 (0.4)
Haliburton, Kawartha, Pine Ridge District Health Unit	2 (0.3)	1 (10.0)	3 (0.4)
Halton Region Public Health	11 (1.6)	0 (0.0)	11 (1.6)
Hastings Prince Edward Public Health	0 (0.0)	0 (0.0)	0 (0.0)
Huron Perth Public Health	0 (0.0)	0 (0.0)	0 (0.0)
Kingston, Frontenac and Lennox & Addington Public Health	2 (0.3)	0 (0.0)	2 (0.3)
Lambton Public Health	1 (0.1)	0 (0.0)	1 (0.1)
Leeds, Grenville & Lanark District Health Unit	2 (0.3)	0 (0.0)	2 (0.3)
Middlesex-London Health Unit	15 (2.2)	0 (0.0)	15 (2.1)
Niagara Region Public Health	8 (1.2)	0 (0.0)	8 (1.1)
North Bay Parry Sound District Health Unit	1 (0.1)	0 (0.0)	1 (0.1)
Northwestern Health Unit	0 (0.0)	0 (0.0)	0 (0.0)
Ottawa Public Health	42 (6.1)	0 (0.0)	42 (6.0)
Peel Public Health	24 (3.5)	0 (0.0)	24 (3.4)
Peterborough Public Health	5 (0.7)	0 (0.0)	5 (0.7)
Porcupine Health Unit	0 (0.0)	0 (0.0)	0 (0.0)

Public Health Unit	Confirmed n (%)	Probable n (%)	Total n (%)
Public Health Sudbury & Districts	1 (0.1)	0 (0.0)	1 (0.1)
Region of Waterloo Public Health and Emergency Services	4 (0.6)	0 (0.0)	4 (0.6)
Renfrew County and District Health Unit	1 (0.1)	0 (0.0)	1 (0.1)
Simcoe Muskoka District Health Unit	3 (0.4)	0 (0.0)	3 (0.4)
Southwestern Public Health	1 (0.2)	0 (0.0)	1 (0.1)
Thunder Bay District Health Unit	0 (0.0)	0 (0.0)	0 (0.0)
Timiskaming Health Unit	0 (0.0)	0 (0.0)	0 (0.0)
Toronto Public Health	505 (73.1)	8 (80.0)	513 (73.2)
Wellington-Dufferin-Guelph Public Health	6 (0.9)	0 (0.0)	6 (0.9)
Windsor-Essex County Health Unit	8 (1.2)	0 (0.0)	8 (1.1)
York Region Public Health	9 (1.3)	0 (0.0)	9 (1.3)
Total	691 (98.6)	10 (1.4)	701 (100.0)

Figure 3. Confirmed MPOX cases and rate per 100,000 population by diagnosing public health unit




Table 4. Severity and outcome of MPOX cases by case classification

Severity and outcome	Confirmed n (%)	Probable n (%)	Total n (%)
Ever hospitalized	20 (2.9)	0 (0.0)	20 (2.9)
Ever in ICU	2 (0.3)	0 (0.0)	2 (0.3)
Outcome: fatal	0 (0.0)	0 (0.0)	0 (0.0)

Risk Factors

- The most commonly reported risk factors include engaging in sexual or intimate contact (e.g., hugging, kissing, cuddling) with new and/or more than one partner.
- Travel, being immunocompromised, or having close contact with someone from outside of the province were less commonly reported.
- Although cases have mostly been identified among males who report sexual or intimate contact with other males (MSM), anyone can get MPOX. Various factors that may increase the potential risk for exposure include close, sexual, and/or other intimate contact with someone who has a MPOX rash, sore, or scabs.

Technical Notes

Data Sources

- The data for this report were based on information entered in the Ontario Ministry of Health (MOH) integrated Public Health Information System (iPHIS) database as of **November 29, 2022 at 1:00 p.m.** and the Public Health Ontario Laboratory Information Management System as of **November 29, 2022 at 9:00 a.m.**
- iPHIS is a dynamic disease reporting system that allows ongoing updates to previously entered data. As a result, data extracted from iPHIS represent a snapshot at the time of extraction and may differ from previous or subsequent reports.

Data Caveats

- These data only represent cases of MPOX reported to public health and recorded in iPHIS. As a result, all case counts are subject to varying degrees of underreporting due to a variety of factors, such as disease awareness and medical care seeking behaviours that may depend on severity of illness, clinical practices, and changes in laboratory testing and reporting behaviours.
- Only MPOX cases meeting the confirmed, probable, or suspect case classification as listed in the [Ontario MOH surveillance case definitions](#) are included in the reported case counts.
- Cases of MPOX are reported based on the Episode Date, which is an estimate of the onset date of disease for a case. In order to determine this date, the following hierarchy exists in iPHIS: Onset Date > Specimen Collection Date > Lab Test Date > Reported Date.
 - For example: If an Onset Date exists, it will be used as the Episode Date. If Onset Date is not available, then the next available date in the hierarchy (i.e., Specimen Collection Date) will be used, and so on.
- Case counts by geography are based on the diagnosing health unit (DHU). DHU refers to the case's public health unit of residence at the time of illness onset or report to public health and not necessarily the location of exposure.
 - Cases for which the DHU was reported as MOHLTC (to signify a case that is not a resident of Ontario) or MUSKOKA-PARRY SOUND (a public health unit that no longer exists) were excluded from this analysis.
- Cases for which the Disposition Status was reported as ENTERED IN ERROR, DOES NOT MEET DEFINITION, DUPLICATE-DO NOT USE, or any variation on these values, were excluded from this analysis.
- The potential for duplicates exists because duplicate sets were not identified and excluded unless they were already resolved at either the local or provincial level prior to data extraction from iPHIS.

Citation

Ontario Agency for Health Protection and Promotion (Public Health Ontario). Epidemiological summary: MPOX in Ontario – May 1, 2022 to November 29, 2022. Toronto, ON: King's Printer for Ontario; 2022.

Disclaimer

This document was developed by Public Health Ontario (PHO). PHO provides scientific and technical advice to Ontario's government, public health organizations and health care providers. PHO's work is guided by the current best available evidence at the time of publication. The application and use of this document is the responsibility of the user. PHO assumes no liability resulting from any such application or use. This document may be reproduced without permission for non-commercial purposes only and provided that appropriate credit is given to PHO. No changes and/or modifications may be made to this document without express written permission from PHO.

Public Health Ontario

Public Health Ontario is an agency of the Government of Ontario dedicated to protecting and promoting the health of all Ontarians and reducing inequities in health. Public Health Ontario links public health practitioners, front-line health workers and researchers to the best scientific intelligence and knowledge from around the world.

For more information about PHO, visit publichealthontario.ca.

©King's Printer for Ontario, 2022

