

WEEKLY EPIDEMIOLOGICAL SUMMARY

SARS-CoV-2 Genomic Surveillance in Ontario,
September 16, 2022

This report summarizes the results of SARS-CoV-2 whole genome sequencing completed by Public Health Ontario as of September 14, 2022 and partner laboratories in the Ontario COVID-19 Genomics Network as of September 13, 2022.

Background

The continued monitoring of global SARS-CoV-2 genomic data has identified changes in the genome as it spreads through populations. These random changes or mutations arise as a virus evolves over time. The accumulation of these mutations can result in a new lineage of the virus, which is a common occurrence. These new lineages will differ slightly in genome sequence and are termed variants. Although many variants will have no difference in the ability to spread or cause disease, some variants have mutations which may enhance virulence, transmissibility, and/or allow the virus to escape natural or vaccine-induced immunity.

The identification of variants and mutations occurs through whole genome sequencing (WGS) of select samples. Through global surveillance of SARS-CoV-2 genomes, a number of variants have been identified with evidence of clinical and/or public health significance, termed variants of concern (VOC). Current VOCs include B.1.1.7 (Alpha), B.1.351 (Beta), P.1 (Gamma), B.1.617.2 (Delta), and B.1.1.529 (Omicron). WGS has also identified a number of variants of interest (VOI), which may share one or more mutations in common with a VOC, but do not have sufficient evidence at this time to be categorized as a VOC (i.e. evidence of increased transmissibility, disease severity, or immune escape). These variants are also characterized and monitored through genomic surveillance. A VOI may be re-classified as a VOC where there is sufficient scientific evidence to support this designation. The VOC/VOI categories used in this report were derived from the [Public Health Agency of Canada \(PHAC\)](#)¹, the [World Health Organization \(WHO\)](#)², and the [European Centre for Disease Prevention and Control \(ECDC\)](#)³.

As SARS-CoV-2 continues to evolve, lineages will naturally divide into descendant sub-lineages – a genetically closely related group derived from a common ancestor. The descendant branches are given new lineage designations, such as the delineation of Omicron to include descendant BA lineages (e.g., BA.1, BA.2, BA.5). The designation of a descendant lineage does not imply a biological difference from the parent lineage. Rather, the new designation represents a refined genetic group that can be tracked separately. As more research is conducted, there may be evidence of a difference, at which time a new WHO Greek letter classification may be assigned to a newly emerged variant.

The Ontario COVID-19 Genomics Network (OCGN) performs WGS on samples received for SARS-CoV-2 diagnostic testing or VOC PCR testing. Sequences are processed using bioinformatics analyses and assigned a Pango lineage⁴ using the pangolin tool⁵, allowing for the identification of VOC, VOI and other lineages.

Highlights

- There were 7,010 cases sequenced by the OCGN for representative surveillance from August 7 to September 3, 2022.
- The proportion of BA.5 cases (including all BA.5 sub-lineages) remained stable at 89.8% (August 21 to 27) and 90.2% (August 28 to September 3) – see Table 5.
- In the most recent week (August 28 to September 3), BA.5.2.1 was the most prevalent lineage (25.6%), followed by BA.5.2 (24.5%), and BA.5.1 (13.8%).
- The proportion of BA.5.2.1 cases remained stable at 25.4% (August 21 to 27) and 25.6% (August 28 to September 3).
 - Based on Nowcast modelling, the proportion of BA.5.2.1 is projected to remain stable reaching 24.6% (95% CI: 22.5% - 26.9%) by September 21, 2022.

Lineage counts may change between reports as the Pango lineage assignment models are updated.

As of December 31, 2021, diagnostic PCR testing was restricted to high-risk populations. Testing eligibility was expanded on April 11, 2022 to include additional high-risk groups (see technical notes for details). As such, representative surveillance only pertains to tested populations.

The OCGN moved from sequencing 20% to 50% of eligible samples on August 24, 2022.

Representative Surveillance


Table 1. Number of COVID-19 cases, number and percentage of cases sequenced for representative surveillance by week, Ontario, August 7 to September 3, 2022

Week	Number of cases	Number sequenced	Percentage sequenced
Week 32 (August 7 - August 13)	9,861	1,484	15.0%
Week 33 (August 14 - August 20)	9,297	1,552	16.7%
Week 34 (August 21 - August 27)	9,045	2,283	25.2%
Week 35 (August 28 - September 3)	7,811	1,691	21.6%
Total	36,014	7,010	19.5%

Note: 'Number of cases' is the number of confirmed positive cases of COVID-19 in Ontario. Date was assigned to best align with sample collection date, which may differ from other PHO products. 'Number sequenced' is the number of cases sequenced for representative surveillance. Results may not be representative of Ontario overall, and do not include all samples tested for other reasons including travel, outbreak investigation, coroner's cases, reinfection or possible vaccine escape. 'Percentage sequenced' may be lower than the sampling proportion because not all cases are eligible to be sequenced (i.e. excludes samples with cycle threshold >30 or insufficient volume). For representative surveillance: details on the proportion of eligible samples sequenced by the OCGN can be found in the technical notes. Week was assigned based on earliest date available for a sample. Results for recent weeks are incomplete as not all sequencing and bioinformatics analyses were complete at the time of data extraction and will be included in subsequent reports.

Data sources: CCM, PHO, Hospital for Sick Children, Kingston Health Sciences Centre, Shared Hospital Laboratory, Hamilton Regional Laboratory Medicine Program

Figure 1. Percentage of COVID-19 cases by the most prevalent lineages and week, representative surveillance, Ontario, September 5, 2021 to September 3, 2022



Note: Results may not be representative of Ontario overall, particularly in earlier weeks. Details on the proportion of eligible samples sequenced by the OCGN can be found in the technical notes. Week was assigned based on earliest date available for a sample. If more than one sample was sequenced for a case, the most recent sample was included. Results for recent weeks are incomplete as not all sequencing and bioinformatics analyses were complete at the time of data extraction and will be included in subsequent reports.

Data sources: PHO, Hospital for Sick Children, Kingston Health Sciences Centre, Shared Hospital Laboratory, Hamilton Regional Laboratory Medicine Program


Table 2. Number and percentage of cases by Pango lineage and week, representative surveillance, Ontario, August 7 to September 3, 2022

WHO label/Pango lineage	Week 32 (August 7 to August 13)	Week 33 (August 14 to August 20)	Week 34 (August 21 to August 27)	Week 35 (August 28 to September 3)	Total (August 7 to September 3)
Omicron	1,484 (100%)	1,552 (100%)	2,283 (100%)	1,691 (100%)	7,010 (100%)
BA.5.2.1	410 (27.6%)	373 (24.0%)	579 (25.4%)	433 (25.6%)	1,795 (25.6%)
BA.5.2	295 (19.9%)	279 (18.0%)	499 (21.9%)	414 (24.5%)	1,487 (21.2%)
BA.5.1	216 (14.6%)	241 (15.5%)	355 (15.5%)	234 (13.8%)	1,046 (14.9%)
BA.4.6	70 (4.7%)	72 (4.6%)	120 (5.3%)	102 (6.0%)	364 (5.2%)
BF.10	83 (5.6%)	110 (7.1%)	97 (4.2%)	68 (4.0%)	358 (5.1%)
BA.5.6	44 (3.0%)	36 (2.3%)	54 (2.4%)	34 (2.0%)	168 (2.4%)
BE.1	25 (1.7%)	59 (3.8%)	49 (2.1%)	30 (1.8%)	163 (2.3%)
BA.5.5	50 (3.4%)	32 (2.1%)	54 (2.4%)	25 (1.5%)	161 (2.3%)
Other Omicron	291 (19.6%)	350 (22.6%)	476 (20.8%)	351 (20.8%)	1,468 (20.9%)
Total sequenced	1,484 (100%)	1,552 (100%)	2,283 (100%)	1,691 (100%)	7,010 (100%)

Note: Includes the most prevalent lineages detected in the past month. Details on the proportion of eligible samples sequenced by the OCGN can be found in the technical notes. Week was assigned based on the earliest date available for the sample. Not all sequencing and bioinformatics analyses for recent weeks were complete at the time of data extraction. Case counts for these weeks may increase in subsequent reports.

Data sources: PHO, Hospital for Sick Children, Kingston Health Sciences Centre, Shared Hospital Laboratory, Hamilton Regional Laboratory Medicine Program

Figure 2. Estimated daily prevalence (%) by Pango lineage, using Nowcast model, Ontario, June 12 to September 24, 2022



Note: Each curve represents the estimated prevalence of a given lineage from Nowcast modelling, which uses 12 weeks of daily representative surveillance data in a multinomial logistic regression. Each set of dots represents the observed daily prevalence of a given lineage, while their size represents the relative number of samples. The vertical dashed line indicates the most recent day of data, after which projected Nowcast prevalence estimates are presented with their 95% confidence intervals. The vertical grey lines indicate the mid-point of the week. Includes all lineages with at least one day of an estimated prevalence of 5% or greater during the 15 week period (12 observed and 3 projected). Only lineages with at least seven days of non-zero case counts were included.

Data sources: PHO, Hospital for Sick Children, Kingston Health Sciences Centre, Shared Hospital Laboratory, Hamilton Regional Laboratory Medicine Program

Table 3. Estimated prevalence (%), projected prevalence (%), and weekly relative growth rate (with 95% confidence intervals) by Pango lineage, using Nowcast model, Ontario, June 12 to September 24, 2022

WHO label/Pango lineage	Week 34 (August 24): Estimated	Week 35 (August 31): Estimated	Week 36 (September 7): Projected	Week 37 (September 14): Projected	Week 38 (September 21): Projected	Weekly relative growth rate
BA.5.2.1	26.8 (25.8 - 27.8)	26.8 (25.7 - 28.0)	26.5 (25.2 - 27.9)	25.9 (24.2 - 27.5)	24.6 (22.5 - 26.9)	1.00 (reference)
BA.5.2	22.1 (21.2 - 23.1)	23.5 (22.3 - 24.6)	24.5 (23.2 - 26.0)	25.3 (23.6 - 27.0)	25.5 (23.1 - 28.0)	1.06 (1.04 - 1.07)
BA.5.1	14.1 (13.3 - 14.9)	13.1 (12.3 - 13.9)	12.0 (11.2 - 12.9)	10.8 (9.9 - 11.8)	9.6 (8.5 - 10.7)	0.93 (0.92 - 0.94)
BA.4.6	5.5 (5.0 - 6.1)	5.9 (5.3 - 6.6)	6.3 (5.5 - 7.2)	6.6 (5.7 - 7.7)	6.8 (5.7 - 8.1)	1.08 (1.05 - 1.10)
BF.10	5.3 (4.8 - 5.8)	5.3 (4.7 - 5.9)	5.3 (4.6 - 6.0)	5.2 (4.4 - 6.0)	4.9 (4.1 - 5.9)	1.00 (0.98 - 1.02)
BA.5.5	2.1 (1.8 - 2.4)	1.7 (1.4 - 1.9)	1.3 (1.1 - 1.6)	1.1 (0.9 - 1.3)	0.8 (0.6 - 1.0)	0.81 (0.79 - 0.83)
BA.2.12.1	0.2 (0.2 - 0.3)	0.1 (0.1 - 0.1)	0.1 (0 - 0.1)	0.0 (0.0 - 0.0)	0.0 (0.0 - 0.0)	0.52 (0.50 - 0.53)
BA.2	0.0 (0.0 - 0.0)	0.0 (0.0 - 0.0)	0.0 (0.0 - 0.0)	0.0 (0.0 - 0.0)	0.0 (0.0 - 0.0)	0.47 (0.45 - 0.49)

Note: The Nowcast model uses 12 weeks of daily representative surveillance data in a multinomial logistic regression that estimates and projects the prevalence of SARS-CoV-2 lineages. The prevalence estimates and projections presented are from the Wednesday (mid-point) of the specified week. Lineages that had at least one day with a prevalence of 5% or greater in the 15 week period (12 observed and 3 projected) were included. Lineages with at least seven days of non-zero case counts were included. Prevalence estimates are based on the model and are not expected to be the same as the observed data (e.g. Table 2). The weekly relative growth rate is a measure of a lineage's growth rate relative to the reference lineage and is estimated in the Nowcast model. Details on the methodology used to calculate Nowcast prevalence estimates, projections, and the weekly relative growth rates can be found in the technical notes.

Data sources: PHO, Hospital for Sick Children, Kingston Health Sciences Centre, Shared Hospital Laboratory, Hamilton Regional Laboratory Medicine Program

Table 4. Number and percentage of cases by Pango lineage and age group, representative surveillance, Ontario, August 7 to September 3, 2022

WHO label/Pango lineage	Ages: 0-4	Ages: 5-11	Ages: 12-19	Ages: 20-39	Ages: 40-59	Ages: 60-79	Ages: 80 and over	Total
Omicron	153 (100%)	66 (100%)	117 (100%)	1,596 (100%)	1,738 (100%)	1,601 (100%)	1,733 (100%)	7,004 (100%)
BA.5.2.1	31 (20.3%)	10 (15.2%)	32 (27.4%)	389 (24.4%)	458 (26.4%)	418 (26.1%)	454 (26.2%)	1,792 (25.6%)
BA.5.2	32 (20.9%)	10 (15.2%)	20 (17.1%)	316 (19.8%)	365 (21.0%)	350 (21.9%)	394 (22.7%)	1,487 (21.2%)
BA.5.1	17 (11.1%)	15 (22.7%)	16 (13.7%)	262 (16.4%)	263 (15.1%)	228 (14.2%)	245 (14.1%)	1,046 (14.9%)
BA.4.6	10 (6.5%)	5 (7.6%)	7 (6.0%)	92 (5.8%)	96 (5.5%)	79 (4.9%)	75 (4.3%)	364 (5.2%)
BF.10	13 (8.5%)	5 (7.6%)	8 (6.8%)	72 (4.5%)	82 (4.7%)	87 (5.4%)	91 (5.3%)	358 (5.1%)
BA.5.6	5 (3.3%)	0 (0.0%)	3 (2.6%)	44 (2.8%)	32 (1.8%)	40 (2.5%)	43 (2.5%)	167 (2.4%)
BE.1	2 (1.3%)	2 (3.0%)	2 (1.7%)	32 (2.0%)	42 (2.4%)	37 (2.3%)	46 (2.7%)	163 (2.3%)
BA.5.5	6 (3.9%)	4 (6.1%)	3 (2.6%)	46 (2.9%)	44 (2.5%)	32 (2.0%)	26 (1.5%)	161 (2.3%)
Other Omicron	37 (24.2%)	15 (22.7%)	26 (22.2%)	343 (21.5%)	356 (20.5%)	330 (20.6%)	359 (20.7%)	1,466 (20.9%)
Total sequenced	153 (100%)	66 (100%)	117 (100%)	1,596 (100%)	1,738 (100%)	1,601 (100%)	1,733 (100%)	7,004 (100%)

Note: Includes the most prevalent lineages detected in the past month. Age was assigned based on the birth date provided in OCGN, and if missing, based on the birth date from CCM; excludes cases with missing birth dates.

Data sources: PHO, Hospital for Sick Children, Kingston Health Sciences Centre, Shared Hospital Laboratory, Hamilton Regional Laboratory Medicine Program, CCM

Table 5. Percentage of BA.5 cases (including all BA.5 sub-lineages), number identified, and total sequenced by public health unit (PHU), region, and week, representative surveillance, Ontario, August 7 to September 3, 2022

Public Health Unit	Week 32 (August 7 to August 13)	Week 33 (August 14 to August 20)	Week 34 (August 21 to August 27)	Week 35 (August 28 to September 3)	Total (August 7 to September 3)
Northwestern Health Unit	100% (2/2)	27.3% (3/11)	63.2% (12/19)	85.7% (6/7)	59.0% (23/39)
Thunder Bay District Health Unit	94.7% (18/19)	92.9% (13/14)	91.7% (22/24)	100% (25/25)	95.1% (78/82)
TOTAL NORTH WEST	95.2% (20/21)	64.0% (16/25)	79.1% (34/43)	96.9% (31/32)	83.5% (101/121)
Algoma Public Health	81.2% (13/16)	75.0% (9/12)	59.3% (16/27)	90.9% (20/22)	75.3% (58/77)
North Bay Parry Sound District Health Unit	90.0% (18/20)	96.2% (25/26)	82.6% (19/23)	25.0% (1/4)	86.3% (63/73)
Porcupine Health Unit	100% (8/8)	83.3% (5/6)	100% (9/9)	95.5% (21/22)	95.6% (43/45)
Public Health Sudbury & Districts	85.3% (29/34)	80.8% (21/26)	93.2% (41/44)	86.5% (45/52)	87.2% (136/156)
Timiskaming Health Unit	0.0% (0/0)	100% (1/1)	50.0% (1/2)	100% (1/1)	75.0% (3/4)
TOTAL NORTH EAST	87.2% (68/78)	85.9% (61/71)	81.9% (86/105)	87.1% (88/101)	85.4% (303/355)
Ottawa Public Health	83.3% (95/114)	84.9% (79/93)	86.3% (126/146)	89.7% (70/78)	85.8% (370/431)
Eastern Ontario Health Unit	78.6% (11/14)	93.8% (15/16)	83.3% (20/24)	100% (13/13)	88.1% (59/67)
Hastings Prince Edward Public Health	73.1% (19/26)	81.6% (31/38)	88.7% (47/53)	95.9% (71/74)	88.0% (168/191)
Kingston, Frontenac and Lennox & Addington Public Health	85.7% (24/28)	90.3% (28/31)	92.0% (80/87)	94.5% (86/91)	92.0% (218/237)
Leeds, Grenville & Lanark District Health Unit	90.0% (18/20)	92.9% (26/28)	100% (17/17)	100% (18/18)	95.2% (79/83)
Renfrew County and District Health Unit	92.9% (13/14)	90.5% (19/21)	97.4% (38/39)	92.0% (23/25)	93.9% (93/99)
TOTAL EASTERN	83.3% (180/216)	87.2% (198/227)	89.6% (328/366)	94.0% (281/299)	89.1% (987/1,108)
Durham Region Health Department	77.8% (21/27)	86.0% (37/43)	75.0% (42/56)	76.1% (51/67)	78.2% (151/193)
Haliburton, Kawartha, Pine Ridge District Health Unit	76.9% (20/26)	86.2% (25/29)	86.8% (33/38)	57.7% (15/26)	78.2% (93/119)
Peel Public Health	90.9% (100/110)	87.4% (90/103)	87.3% (137/157)	87.9% (102/116)	88.3% (429/486)
Peterborough Public Health	100% (4/4)	100% (5/5)	100% (12/12)	100% (7/7)	100% (28/28)
Simcoe Muskoka District Health Unit	80.8% (42/52)	92.8% (64/69)	88.5% (77/87)	88.9% (72/81)	88.2% (255/289)
York Region Public Health	88.7% (141/159)	89.3% (134/150)	95.0% (151/159)	92.9% (92/99)	91.4% (518/567)
TOTAL CENTRAL EAST	86.8% (328/378)	89.0% (355/399)	88.8% (452/509)	85.6% (339/396)	87.6% (1,474/1,682)

Public Health Unit	Week 32 (August 7 to August 13)	Week 33 (August 14 to August 20)	Week 34 (August 21 to August 27)	Week 35 (August 28 to September 3)	Total (August 7 to September 3)
Toronto Public Health	91.7% (297/324)	92.4% (304/329)	89.3% (308/345)	90.3% (318/352)	90.9% (1,227/1,350)
TOTAL TORONTO	91.7% (297/324)	92.4% (304/329)	89.3% (308/345)	90.3% (318/352)	90.9% (1,227/1,350)
Chatham-Kent Public Health	100% (11/11)	100% (17/17)	96.2% (25/26)	93.3% (14/15)	97.1% (67/69)
Grey Bruce Health Unit	100% (16/16)	100% (13/13)	90.0% (27/30)	70.6% (12/17)	89.5% (68/76)
Huron Perth Public Health	88.9% (8/9)	80.0% (8/10)	92.0% (23/25)	81.8% (9/11)	87.3% (48/55)
Lambton Public Health	91.7% (11/12)	88.9% (16/18)	85.2% (23/27)	90.9% (10/11)	88.2% (60/68)
Middlesex-London Health Unit	89.8% (53/59)	95.9% (70/73)	95.6% (131/137)	97.9% (92/94)	95.3% (346/363)
Southwestern Public Health	100% (17/17)	92.9% (26/28)	88.2% (45/51)	90.9% (50/55)	91.4% (138/151)
Windsor-Essex County Health Unit	88.6% (31/35)	96.7% (29/30)	94.3% (83/88)	98.0% (48/49)	94.6% (191/202)
TOTAL SOUTH WEST	92.5% (147/159)	94.7% (179/189)	93.0% (357/384)	93.3% (235/252)	93.3% (918/984)
Brant County Health Unit	84.0% (21/25)	97.6% (41/42)	72.7% (16/22)	100% (6/6)	88.4% (84/95)
City of Hamilton Public Health Services	90.1% (91/101)	90.7% (88/97)	92.6% (176/190)	91.0% (71/78)	91.4% (426/466)
Haldimand-Norfolk Health Unit	84.6% (11/13)	100% (9/9)	71.4% (10/14)	94.4% (17/18)	87% (47/54)
Halton Region Public Health	94.3% (33/35)	88.6% (31/35)	92.3% (48/52)	84.8% (28/33)	90.3% (140/155)
Niagara Region Public Health	97.1% (33/34)	86.1% (31/36)	97.6% (80/82)	83.0% (44/53)	91.7% (188/205)
Region of Waterloo Public Health and Emergency Services	85.9% (55/64)	91.0% (61/67)	90.5% (114/126)	92.5% (37/40)	89.9% (267/297)
Wellington-Dufferin-Guelph Public Health	91.7% (33/36)	79.2% (19/24)	90.9% (40/44)	96.8% (30/31)	90.4% (122/135)
TOTAL CENTRAL WEST	89.9% (277/308)	90.3% (280/310)	91.3% (484/530)	90.0% (233/259)	90.5% (1,274/1,407)
UNKNOWN	0.0% (0/0)	100% (2/2)	100% (1/1)	0.0% (0/0)	100% (3/3)
TOTAL ONTARIO	88.7% (1,317/1,484)	89.9% (1,395/1,552)	89.8% (2,050/2,283)	90.2% (1,525/1,691)	89.7% (6,287/7,010)

Note: Details on the proportion of eligible samples sequenced by the OCGN can be found in the technical notes. Week was assigned based on the earliest date available for the sample. Not all sequencing and bioinformatics analyses for recent weeks were complete at the time of data extraction. Case counts for these weeks may increase in subsequent reports. Public health unit was assigned based on diagnosing health unit in CCM. If a case did not link to CCM (2.9%), OCGN patient postal code was used. Ordering provider postal code was used if patient postal code was missing.

Data sources: PHO, Hospital for Sick Children, Kingston Health Sciences Centre, Shared Hospital Laboratory, Hamilton Regional Laboratory Medicine Program

Table 6. Number and percentage (row %) of cases by vaccine category and Pango lineage, representative surveillance, Ontario, August 7 to September 3, 2022

WHO label / Pango lineage	Unvaccinated	Post-series initiation	Post-series completion	Post-booster dose	Post-two booster doses	Total cases
Omicron	1,264 (18.6%)	55 (0.8%)	728 (10.7%)	2,774 (40.8%)	1,983 (29.1%)	6,804 (100%)
BA.5.2.1	305 (17.5%)	9 (0.5%)	174 (10.0%)	720 (41.3%)	536 (30.7%)	1,744 (100%)
BA.5.2	263 (18.1%)	16 (1.1%)	154 (10.6%)	591 (40.7%)	427 (29.4%)	1,451 (100%)
BA.5.1	219 (21.5%)	9 (0.9%)	116 (11.4%)	387 (37.9%)	289 (28.3%)	1,020 (100%)
BA.4.6	63 (17.9%)	4 (1.1%)	30 (8.5%)	155 (44.2%)	99 (28.2%)	351 (100%)
BF.10	73 (20.7%)	6 (1.7%)	31 (8.8%)	142 (40.3%)	100 (28.4%)	352 (100%)
BA.5.6	27 (17.1%)	0 (0.0%)	16 (10.1%)	71 (44.9%)	44 (27.8%)	158 (100%)
BE.1	25 (16.2%)	1 (0.6%)	16 (10.4%)	54 (35.1%)	58 (37.7%)	154 (100%)
BA.5.5	24 (15.5%)	1 (0.6%)	29 (18.7%)	68 (43.9%)	33 (21.3%)	155 (100%)
Other Omicron	265 (18.7%)	9 (0.6%)	162 (11.4%)	586 (41.3%)	397 (28.0%)	1,419 (100%)
Total sequenced	1,264 (18.6%)	55 (0.8%)	728 (10.7%)	2,774 (40.8%)	1,983 (29.1%)	6,804 (100%)

Note: Includes the most prevalent lineages detected in the past month. Cases include those that linked to CCM (97.1%). Individuals with a vaccine not approved by Health Canada were excluded. Vaccine category definitions can be found in the [Confirmed Cases of COVID-19 Following Vaccination in Ontario](#) report. A higher proportion of cases reported among more vaccinated cases is a reflection of both trends in vaccine administration (increasing number of doses administered over time) and trends in COVID-19 incidence. Details on the proportion of eligible samples sequenced by the OCGN can be found in the technical notes. Sample date represents the earliest date available for the sample. Not all sequencing and bioinformatics analyses for recent weeks were complete at the time of data extraction.

Data sources: PHO, Hospital for Sick Children, Kingston Health Sciences Centre, Hamilton Regional Laboratory Medicine Program, Shared Hospital Laboratory, CCM, COVaxON

Table 7a. Percentage of cases ever admitted to hospital by vaccination status and Pango lineage, representative surveillance, Ontario, August 7 to September 3, 2022

WHO label / Pango lineage	Unvaccinated	Post-series initiation	Post-series completion	Post-booster dose	Post-two booster doses
Omicron	4.4% (56/1,264)	9.1% (5/55)	6.7% (49/728)	5.0% (140/2,774)	6.3% (125/1,983)
BA.5.2.1	5.2% (16/305)	11.1% (1/9)	5.2% (9/174)	5.0% (36/720)	5.6% (30/536)
BA.5.2	4.2% (11/263)	12.5% (2/16)	5.8% (9/154)	4.6% (27/591)	6.1% (26/427)
BA.5.1	4.1% (9/219)	11.1% (1/9)	7.8% (9/116)	4.7% (18/387)	6.2% (18/289)
BA.4.6	4.8% (3/63)	0.0% (0/4)	6.7% (2/30)	3.9% (6/155)	6.1% (6/99)
BF.10	6.8% (5/73)	16.7% (1/6)	16.1% (5/31)	6.3% (9/142)	5.0% (5/100)
BA.5.6	0.0% (0/27)	0.0% (0/0)	6.2% (1/16)	11.3% (8/71)	4.5% (2/44)
BE.1	8.0% (2/25)	0.0% (0/1)	12.5% (2/16)	3.7% (2/54)	5.2% (3/58)
BA.5.5	8.3% (2/24)	0.0% (0/1)	0.0% (0/29)	8.8% (6/68)	3.0% (1/33)
Other Omicron	3.0% (8/265)	0.0% (0/9)	7.4% (12/162)	4.8% (28/586)	8.6% (34/397)
Total sequenced	4.4% (56/1,264)	9.1% (5/55)	6.7% (49/728)	5.0% (140/2,774)	6.3% (125/1,983)

Note: Includes the most prevalent lineages detected in the past month. Percentage is the number of cases ever admitted to hospital (at the time of data extraction) divided by the total number of cases in that lineage and vaccine category. Cases include only those that linked to CCM (97.1%). Individuals with a vaccine not approved by Health Canada were excluded. Vaccine category definitions can be found in the [Confirmed Cases of COVID-19 Following Vaccination in Ontario](#) report. Factors, such as age, that may affect the risk of COVID-19 hospital admission and death are not accounted for in these analyses. As a result, the proportion of severe outcomes following booster doses may be higher compared to other vaccination statuses since booster eligibility was initially restricted to older adults who are more likely to have severe outcomes and who also have the highest uptake for booster doses. Results may not be representative of Ontario overall. Details on the proportion of eligible samples sequenced by the OCGN can be found in the technical notes. Sample date represents the earliest date available for the sample. Not all sequencing and bioinformatics analyses for recent weeks were complete at the time of data extraction.

Data sources: PHO, Hospital for Sick Children, Kingston Health Sciences Centre, Hamilton Regional Laboratory Medicine Program, The Shared Hospital Laboratory, CCM (hospital admission indicator), COVaxON

Table 7b. Percentage of deceased cases by vaccination status and Pango lineage, representative surveillance, Ontario, August 7 to September 3, 2022

WHO label / Pango lineage	Unvaccinated	Post-series initiation	Post-series completion	Post-booster dose	Post-two booster doses
Omicron	0.6% (7/1,264)	3.6% (2/55)	0.7% (5/728)	0.5% (15/2,774)	1.8% (35/1,983)
BA.5.2.1	0.3% (1/305)	0.0% (0/9)	0.6% (1/174)	0.7% (5/720)	0.9% (5/536)
BA.5.2	0.4% (1/263)	6.2% (1/16)	0.6% (1/154)	0.0% (0/591)	1.6% (7/427)
BA.5.1	0.9% (2/219)	0.0% (0/9)	0.0% (0/116)	0.5% (2/387)	1.7% (5/289)
BA.4.6	3.2% (2/63)	0.0% (0/4)	0.0% (0/30)	0.6% (1/155)	1.0% (1/99)
BF.10	0.0% (0/73)	0.0% (0/6)	0.0% (0/31)	0.7% (1/142)	3.0% (3/100)
BA.5.6	0.0% (0/27)	0.0% (0/0)	0.0% (0/16)	1.4% (1/71)	2.3% (1/44)
BE.1	0.0% (0/25)	0.0% (0/1)	6.2% (1/16)	1.9% (1/54)	1.7% (1/58)
BA.5.5	0.0% (0/24)	0.0% (0/1)	0.0% (0/29)	0.0% (0/68)	0.0% (0/33)
Other Omicron	0.4% (1/265)	11.1% (1/9)	1.2% (2/162)	0.7% (4/586)	3.0% (12/397)
Total sequenced	0.6% (7/1,264)	3.6% (2/55)	0.7% (5/728)	0.5% (15/2,774)	1.8% (35/1,983)

Note: Includes the most prevalent lineages detected in the past month. Percentage is the number of deceased cases divided by the total number of cases in that lineage and vaccine category. Cases include only those that linked to CCM (97.1%). Individuals with a vaccine not approved by Health Canada were excluded. Vaccine category definitions can be found in the [Confirmed Cases of COVID-19 Following Vaccination in Ontario](#) report. Deceased cases include cases that reported a “Fatal” outcome and the type of death value in CCM was not ‘DOPHS was unrelated to cause of death’ or ‘Under PHU Review’ at the time of data extraction. Factors, such as age, that may affect the risk of COVID-19 hospitalization and death are not accounted for in these analyses. As a result, the proportion of severe outcomes following booster doses may be higher compared to other vaccination statuses since booster eligibility was initially restricted to older adults who are more likely to have severe outcomes and who also have the highest uptake for booster doses. Results may not be representative of Ontario overall. Details on the proportion of eligible samples sequenced by the OCGN can be found in the technical notes. Sample date represents the earliest date available for the sample. Not all sequencing and bioinformatics analyses for recent weeks were complete at the time of data extraction.

Data sources: PHO, Hospital for Sick Children, Kingston Health Sciences Centre, Hamilton Regional Laboratory Medicine Program, The Shared Hospital Laboratory, CCM (deceased indicator), COVaxON

Table 8a. Number and percentage of cases by Pango lineage and public health unit (PHU), representative surveillance, North West Region, August 7 to September 3, 2022

WHO label / Pango lineage	Northwestern Health Unit	Thunder Bay District Health Unit	Total
Omicron	39 (100%)	82 (100%)	121 (100%)
BA.5.2.1	9 (23.1%)	22 (26.8%)	31 (25.6%)
BA.5.2	6 (15.4%)	30 (36.6%)	36 (29.8%)
BA.5.1	3 (7.7%)	5 (6.1%)	8 (6.6%)
BA.4.6	1 (2.6%)	2 (2.4%)	3 (2.5%)
BF.10	2 (5.1%)	1 (1.2%)	3 (2.5%)
BA.5.6	0 (0.0%)	0 (0.0%)	0 (0.0%)
BE.1	0 (0.0%)	7 (8.5%)	7 (5.8%)
BA.5.5	0 (0.0%)	2 (2.4%)	2 (1.7%)
Other Omicron	18 (46.2%)	13 (15.9%)	31 (25.6%)
Total sequenced	39 (100%)	82 (100%)	121 (100%)

Note: Includes the most prevalent lineages detected in the past month. Details on the proportion of eligible samples sequenced by the OCGN can be found in the technical notes. Sample date represents the earliest date available for the sample. Not all sequencing and bioinformatics analyses for recent weeks were complete at the time of data extraction. Public health unit was assigned based on diagnosing health unit in CCM. If a case did not link to CCM (2.9%), OCGN patient postal code was used. Ordering provider postal code was used if patient postal code was missing.

Data sources: PHO, Hospital for Sick Children, Kingston Health Sciences Centre, Shared Hospital Laboratory, Hamilton Regional Laboratory Medicine Program, CCM

Table 8b. Number and percentage of cases by Pango lineage and public health unit (PHU), representative surveillance, North East Region, August 7 to September 3, 2022

WHO label / Pango lineage	Algoma Public Health	North Bay Parry Sound District Health Unit	Porcupine Health Unit	Public Health Sudbury & Districts	Timiskaming Health Unit	Total
Omicron	77 (100%)	73 (100%)	45 (100%)	156 (100%)	4 (100%)	355 (100%)
BA.5.2.1	20 (26.0%)	15 (20.5%)	10 (22.2%)	39 (25.0%)	0 (0.0%)	84 (23.7%)
BA.5.2	14 (18.2%)	10 (13.7%)	3 (6.7%)	59 (37.8%)	3 (75.0%)	89 (25.1%)
BA.5.1	8 (10.4%)	32 (43.8%)	14 (31.1%)	9 (5.8%)	0 (0.0%)	63 (17.7%)
BA.4.6	15 (19.5%)	7 (9.6%)	1 (2.2%)	14 (9.0%)	0 (0.0%)	37 (10.4%)
BF.10	0 (0.0%)	1 (1.4%)	9 (20.0%)	4 (2.6%)	0 (0.0%)	14 (3.9%)
BA.5.6	2 (2.6%)	0 (0.0%)	0 (0.0%)	1 (0.6%)	0 (0.0%)	3 (0.8%)
BE.1	0 (0.0%)	3 (4.1%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	3 (0.8%)
BA.5.5	1 (1.3%)	0 (0.0%)	0 (0.0%)	2 (1.3%)	0 (0.0%)	3 (0.8%)
Other Omicron	17 (22.1%)	5 (6.8%)	8 (17.8%)	28 (17.9%)	1 (25.0%)	59 (16.6%)
Total sequenced	77 (100%)	73 (100%)	45 (100%)	156 (100%)	4 (100%)	355 (100%)

Note: Includes the most prevalent lineages detected in the past month. Details on the proportion of eligible samples sequenced by the OCGN can be found in the technical notes. Sample date represents the earliest date available for the sample. Not all sequencing and bioinformatics analyses for recent weeks were complete at the time of data extraction. Public health unit was assigned based on diagnosing health unit in CCM. If a case did not link to CCM (2.9%), OCGN patient postal code was used. Ordering provider postal code was used if patient postal code was missing.

Data sources: PHO, Hospital for Sick Children, Kingston Health Sciences Centre, Shared Hospital Laboratory, Hamilton Regional Laboratory Medicine Program, CCM

Table 8c. Number and percentage of cases by Pango lineage and public health unit (PHU), representative surveillance, Eastern Region, August 7 to September 3, 2022

WHO label / Pango lineage	Eastern Ontario Health Unit	Hastings Prince Edward Public Health	Kingston, Frontenac and Lennox & Addington Public Health	Leeds, Grenville & Lanark District Health Unit	Ottawa Public Health	Renfrew County and District Health Unit	Total
Omicron	67 (100%)	191 (100%)	237 (100%)	83 (100%)	431 (100%)	99 (100%)	1,108 (100%)
BA.5.2.1	19 (28.4%)	41 (21.5%)	124 (52.3%)	21 (25.3%)	99 (23.0%)	5 (5.1%)	309 (27.9%)
BA.5.2	11 (16.4%)	24 (12.6%)	40 (16.9%)	9 (10.8%)	76 (17.6%)	45 (45.5%)	205 (18.5%)
BA.5.1	9 (13.4%)	13 (6.8%)	22 (9.3%)	10 (12.0%)	73 (16.9%)	3 (3.0%)	130 (11.7%)
BA.4.6	3 (4.5%)	8 (4.2%)	10 (4.2%)	1 (1.2%)	39 (9.0%)	5 (5.1%)	66 (6.0%)
BF.10	1 (1.5%)	5 (2.6%)	8 (3.4%)	19 (22.9%)	42 (9.7%)	25 (25.3%)	100 (9.0%)
BA.5.6	1 (1.5%)	0 (0.0%)	1 (0.4%)	2 (2.4%)	13 (3.0%)	1 (1.0%)	18 (1.6%)
BE.1	2 (3.0%)	13 (6.8%)	1 (0.4%)	0 (0.0%)	6 (1.4%)	1 (1.0%)	23 (2.1%)
BA.5.5	0 (0.0%)	1 (0.5%)	3 (1.3%)	1 (1.2%)	3 (0.7%)	3 (3.0%)	11 (1.0%)
Other Omicron	21 (31.3%)	86 (45.0%)	28 (11.8%)	20 (24.1%)	80 (18.6%)	11 (11.1%)	246 (22.2%)
Total sequenced	67 (100%)	191 (100%)	237 (100%)	83 (100%)	431 (100%)	99 (100%)	1,108 (100%)

Note: Includes the most prevalent lineages detected in the past month. Details on the proportion of eligible samples sequenced by the OCGN can be found in the technical notes. Sample date represents the earliest date available for the sample. Not all sequencing and bioinformatics analyses for recent weeks were complete at the time of data extraction. Public health unit was assigned based on diagnosing health unit in CCM. If a case did not link to CCM (2.9%), OCGN patient postal code was used. Ordering provider postal code was used if patient postal code was missing.

Data sources: PHO, Hospital for Sick Children, Kingston Health Sciences Centre, Shared Hospital Laboratory, Hamilton Regional Laboratory Medicine Program, CCM

Table 8d. Number and percentage of cases by Pango lineage and public health unit (PHU), representative surveillance, Central East Region, August 7 to September 3, 2022

WHO label / Pango lineage	Durham Region Health Department	Haliburton, Kawartha, Pine Ridge District Health Unit	Peel Public Health	Peterborough Public Health	Simcoe Muskoka District Health Unit	York Region Public Health	Total
Omicron	193 (100%)	119 (100%)	486 (100%)	28 (100%)	289 (100%)	567 (100%)	1,682 (100%)
BA.5.2.1	44 (22.8%)	28 (23.5%)	135 (27.8%)	3 (10.7%)	72 (24.9%)	123 (21.7%)	405 (24.1%)
BA.5.2	36 (18.7%)	15 (12.6%)	95 (19.5%)	5 (17.9%)	42 (14.5%)	141 (24.9%)	334 (19.9%)
BA.5.1	23 (11.9%)	11 (9.2%)	59 (12.1%)	1 (3.6%)	65 (22.5%)	78 (13.8%)	237 (14.1%)
BA.4.6	28 (14.5%)	23 (19.3%)	23 (4.7%)	0 (0.0%)	22 (7.6%)	25 (4.4%)	121 (7.2%)
BF.10	11 (5.7%)	20 (16.8%)	17 (3.5%)	2 (7.1%)	17 (5.9%)	34 (6.0%)	101 (6.0%)
BA.5.6	2 (1.0%)	0 (0.0%)	20 (4.1%)	5 (17.9%)	7 (2.4%)	26 (4.6%)	60 (3.6%)
BE.1	4 (2.1%)	1 (0.8%)	8 (1.6%)	1 (3.6%)	3 (1.0%)	10 (1.8%)	27 (1.6%)
BA.5.5	2 (1.0%)	3 (2.5%)	7 (1.4%)	1 (3.6%)	7 (2.4%)	24 (4.2%)	44 (2.6%)
Other Omicron	43 (22.3%)	18 (15.1%)	122 (25.1%)	10 (35.7%)	54 (18.7%)	106 (18.7%)	353 (21.0%)
Total sequenced	193 (100%)	119 (100%)	486 (100%)	28 (100%)	289 (100%)	567 (100%)	1,682 (100%)

Note: Includes the most prevalent lineages detected in the past month. Details on the proportion of eligible samples sequenced by the OCGN can be found in the technical notes. Sample date represents the earliest date available for the sample. Not all sequencing and bioinformatics analyses for recent weeks were complete at the time of data extraction. Public health unit was assigned based on diagnosing health unit in CCM. If a case did not link to CCM (2.9%), OCGN patient postal code was used. Ordering provider postal code was used if patient postal code was missing.

Data sources: PHO, Hospital for Sick Children, Kingston Health Sciences Centre, Shared Hospital Laboratory, Hamilton Regional Laboratory Medicine Program, CCM

Table 8e. Number and percentage of cases by Pango lineage and public health unit (PHU), representative surveillance, Toronto Region, August 7 to September 3, 2022

WHO label / Pango lineage	Toronto Public Health	Total
Omicron	1,350 (100%)	1,350 (100%)
BA.5.2.1	363 (26.9%)	363 (26.9%)
BA.5.2	297 (22.0%)	297 (22.0%)
BA.5.1	226 (16.7%)	226 (16.7%)
BA.4.6	51 (3.8%)	51 (3.8%)
BF.10	48 (3.6%)	48 (3.6%)
BA.5.6	29 (2.1%)	29 (2.1%)
BE.1	25 (1.9%)	25 (1.9%)
BA.5.5	37 (2.7%)	37 (2.7%)
Other Omicron	274 (20.3%)	274 (20.3%)
Total sequenced	1,350 (100%)	1,350 (100%)

Note: Includes the most prevalent lineages detected in the past month. Details on the proportion of eligible samples sequenced by the OCGN can be found in the technical notes. Sample date represents the earliest date available for the sample. Not all sequencing and bioinformatics analyses for recent weeks were complete at the time of data extraction. Public health unit was assigned based on diagnosing health unit in CCM. If a case did not link to CCM (2.9%), OCGN patient postal code was used. Ordering provider postal code was used if patient postal code was missing.

Data sources: PHO, Hospital for Sick Children, Kingston Health Sciences Centre, Shared Hospital Laboratory, Hamilton Regional Laboratory Medicine Program, CCM

Table 8f. Number and percentage of cases by Pango lineage and public health unit (PHU), representative surveillance, South West Region, August 7 to September 3, 2022

WHO label / Pango lineage	Chatham-Kent Public Health	Grey Bruce Health Unit	Huron Perth Public Health	Lambton Public Health	Middlesex-London Health Unit	Southwestern Public Health	Windsor-Essex County Health Unit	Total
Omicron	69 (100%)	76 (100%)	55 (100%)	68 (100%)	363 (100%)	151 (100%)	202 (100%)	984 (100%)
BA.5.2.1	16 (23.2%)	23 (30.3%)	18 (32.7%)	17 (25.0%)	96 (26.4%)	29 (19.2%)	52 (25.7%)	251 (25.5%)
BA.5.2	6 (8.7%)	10 (13.2%)	10 (18.2%)	15 (22.1%)	65 (17.9%)	56 (37.1%)	44 (21.8%)	206 (20.9%)
BA.5.1	4 (5.8%)	16 (21.1%)	4 (7.3%)	10 (14.7%)	83 (22.9%)	15 (9.9%)	53 (26.2%)	185 (18.8%)
BA.4.6	0 (0.0%)	1 (1.3%)	4 (7.3%)	1 (1.5%)	5 (1.4%)	9 (6.0%)	4 (2.0%)	24 (2.4%)
BF.10	6 (8.7%)	2 (2.6%)	3 (5.5%)	1 (1.5%)	6 (1.7%)	1 (0.7%)	8 (4.0%)	27 (2.7%)
BA.5.6	0 (0.0%)	1 (1.3%)	2 (3.6%)	1 (1.5%)	7 (1.9%)	5 (3.3%)	4 (2.0%)	20 (2.0%)
BE.1	1 (1.4%)	2 (2.6%)	2 (3.6%)	0 (0.0%)	46 (12.7%)	4 (2.6%)	1 (0.5%)	56 (5.7%)
BA.5.5	4 (5.8%)	8 (10.5%)	2 (3.6%)	1 (1.5%)	5 (1.4%)	0 (0.0%)	8 (4.0%)	28 (2.8%)
Other Omicron	32 (46.4%)	13 (17.1%)	10 (18.2%)	22 (32.4%)	50 (13.8%)	32 (21.2%)	28 (13.9%)	187 (19.0%)
Total sequenced	69 (100%)	76 (100%)	55 (100%)	68 (100%)	363 (100%)	151 (100%)	202 (100%)	984 (100%)

Note: Includes the most prevalent lineages detected in the past month. Details on the proportion of eligible samples sequenced by the OCGN can be found in the technical notes. Sample date represents the earliest date available for the sample. Not all sequencing and bioinformatics analyses for recent weeks were complete at the time of data extraction. Public health unit was assigned based on diagnosing health unit in CCM. If a case did not link to CCM (2.9%), OCGN patient postal code was used. Ordering provider postal code was used if patient postal code was missing.

Data sources: PHO, Hospital for Sick Children, Kingston Health Sciences Centre, Shared Hospital Laboratory, Hamilton Regional Laboratory Medicine Program, CCM

Table 8g. Number and percentage of cases by Pango lineage and public health unit (PHU), representative surveillance, Central West Region, August 7 to September 3, 2022

WHO label / Pango lineage	Brant County Health Unit	City of Hamilton Public Health Services	Haldimand- Norfolk Health Unit	Halton Region Public Health	Niagara Region Public Health	Region of Waterloo Public Health and Emergency Services	Wellington- Dufferin- Guelph Public Health	Total
Omicron	95 (100%)	466 (100%)	54 (100%)	155 (100%)	205 (100%)	297 (100%)	135 (100%)	1,407 (100%)
BA.5.2.1	30 (31.6%)	138 (29.6%)	19 (35.2%)	30 (19.4%)	53 (25.9%)	53 (17.8%)	28 (20.7%)	351 (24.9%)
BA.5.2	18 (18.9%)	96 (20.6%)	10 (18.5%)	37 (23.9%)	56 (27.3%)	73 (24.6%)	30 (22.2%)	320 (22.7%)
BA.5.1	16 (16.8%)	57 (12.2%)	6 (11.1%)	27 (17.4%)	29 (14.1%)	35 (11.8%)	26 (19.3%)	196 (13.9%)
BA.4.6	1 (1.1%)	19 (4.1%)	5 (9.3%)	12 (7.7%)	2 (1.0%)	14 (4.7%)	9 (6.7%)	62 (4.4%)
BF.10	6 (6.3%)	15 (3.2%)	0 (0.0%)	6 (3.9%)	15 (7.3%)	21 (7.1%)	2 (1.5%)	65 (4.6%)
BA.5.6	2 (2.1%)	18 (3.9%)	1 (1.9%)	4 (2.6%)	3 (1.5%)	6 (2.0%)	3 (2.2%)	37 (2.6%)
BE.1	0 (0.0%)	8 (1.7%)	0 (0.0%)	1 (0.6%)	7 (3.4%)	3 (1.0%)	3 (2.2%)	22 (1.6%)
BA.5.5	1 (1.1%)	12 (2.6%)	1 (1.9%)	8 (5.2%)	3 (1.5%)	8 (2.7%)	3 (2.2%)	36 (2.6%)
Other Omicron	21 (22.1%)	103 (22.1%)	12 (22.2%)	30 (19.4%)	37 (18.0%)	84 (28.3%)	31 (23.0%)	318 (22.6%)
Total sequenced	95 (100%)	466 (100%)	54 (100%)	155 (100%)	205 (100%)	297 (100%)	135 (100%)	1,407 (100%)

Note: Includes the most prevalent lineages detected in the past month. Details on the proportion of eligible samples sequenced by the OCGN can be found in the technical notes. Sample date represents the earliest date available for the sample. Not all sequencing and bioinformatics analyses for recent weeks were complete at the time of data extraction. Public health unit was assigned based on diagnosing health unit in CCM. If a case did not link to CCM (2.9%), OCGN patient postal code was used. Ordering provider postal code was used if patient postal code was missing.

Data sources: PHO, Hospital for Sick Children, Kingston Health Sciences Centre, Shared Hospital Laboratory, Hamilton Regional Laboratory Medicine Program, CCM

Table 9. Number and percentage (row %) of outbreak-associated and non outbreak-associated cases by Pango lineage, representative surveillance, Ontario, August 7 to September 3, 2022

WHO label / Pango lineage	Outbreak-associated	Non outbreak-associated	Total cases
Omicron	2,412 (35.4%)	4,396 (64.6%)	6,808 (100%)
BA.5.2.1	682 (39.1%)	1,063 (60.9%)	1,745 (100%)
BA.5.2	552 (38.0%)	900 (62.0%)	1,452 (100%)
BA.5.1	350 (34.3%)	670 (65.7%)	1,020 (100%)
BA.4.6	99 (28.2%)	252 (71.8%)	351 (100%)
BF.10	132 (37.5%)	220 (62.5%)	352 (100%)
BA.5.6	44 (27.8%)	114 (72.2%)	158 (100%)
BE.1	70 (45.2%)	85 (54.8%)	155 (100%)
BA.5.5	26 (16.7%)	130 (83.3%)	156 (100%)
Other Omicron	457 (32.2%)	962 (67.8%)	1,419 (100%)
Total sequenced	2,412 (35.4%)	4,396 (64.6%)	6,808 (100%)

Note: Includes the most prevalent lineages detected in the past month. Cases include only those that linked to CCM (97.1%). ‘Outbreak-associated cases’ include cases linked to a confirmed outbreak as declared by the local medical officer of health or their designate. Details on the proportion of eligible samples sequenced by the OCGN can be found in the technical notes. Sample date represents the earliest date available for the sample. Not all sequencing and bioinformatics analyses for recent weeks were complete at the time of data extraction.

Data sources: PHO, Hospital for Sick Children, Kingston Health Sciences Centre, Hamilton Regional Laboratory Medicine Program, The Shared Hospital Laboratory, CCM (outbreak Indicator)

Cumulative Whole Genome Sequencing Results

Table 10. Number of cases by Pango lineage, cumulative counts, Ontario, January 1, 2021 to September 3, 2022

WHO label / Pango lineage	January 1, 2021 – August 6, 2022	August 7 – September 3, 2022	Total
Variant of Concern (VOC)			
Alpha	11,807	0	11,807
Beta	1,230	0	1,230
Gamma	3,927	0	3,927
Delta	46,043	0	46,043
Omicron	86,116	7,781	93,897
B.1.1.529	27	0	27
BA.1	6,399	1	6,400
BA.1.1	18,889	0	18,889
BA.1.1.1	44	0	44
BA.1.1.10	1,693	0	1,693
BA.1.1.11	1	0	1
BA.1.1.12	1	0	1
BA.1.1.13	5	0	5
BA.1.1.14	522	0	522
BA.1.1.15	3	0	3
BA.1.1.16	99	0	99
BA.1.1.17	6	0	6
BA.1.1.18	134	0	134
BA.1.1.2	6	0	6
BA.1.1.6	3	0	3
BA.1.1.7	5	0	5
BA.1.1.8	3	0	3
BA.1.12	1	0	1
BA.1.13	35	0	35
BA.1.13.1	1	0	1
BA.1.14	70	0	70
BA.1.14.1	1	0	1
BA.1.14.2	1	0	1
BA.1.15	1,653	0	1,653
BA.1.15.1	22	0	22
BA.1.15.2	3	0	3
BA.1.16	2	0	2
BA.1.17	711	1	712
BA.1.17.2	1,533	0	1,533

WHO label / Pango lineage	January 1, 2021 – August 6, 2022	August 7 – September 3, 2022	Total
BA.1.18	93	0	93
BA.1.19	23	0	23
BA.1.20	91	0	91
BA.1.21	8	0	8
BA.1.3	1	0	1
BA.1.6	66	0	66
BA.1.7	1	0	1
BA.1.9	37	0	37
BA.2	15,336	5	15,341
BA.2.1	137	0	137
BA.2.10	389	0	389
BA.2.10.1	16	0	16
BA.2.10.3	1	0	1
BA.2.11	7	0	7
BA.2.12	63	0	63
BA.2.12.1	6,445	43	6,488
BA.2.12.2	2	0	2
BA.2.13	26	0	26
BA.2.13.1	44	1	45
BA.2.15	1	0	1
BA.2.16	8	0	8
BA.2.17	29	0	29
BA.2.18	68	2	70
BA.2.2	4	0	4
BA.2.20	1,688	0	1,688
BA.2.21	328	0	328
BA.2.22	5	0	5
BA.2.23	59	0	59
BA.2.24	2	0	2
BA.2.25	1	0	1
BA.2.26	8	0	8
BA.2.27	1	0	1
BA.2.3	2,369	0	2,369
BA.2.3.10	79	0	79
BA.2.3.12	3	0	3
BA.2.3.14	2	0	2
BA.2.3.15	1	0	1
BA.2.3.16	5	0	5
BA.2.3.17	3	0	3
BA.2.3.2	21	0	21

WHO label / Pango lineage	January 1, 2021 – August 6, 2022	August 7 – September 3, 2022	Total
BA.2.3.4	957	0	957
BA.2.3.5	1	0	1
BA.2.3.6	79	0	79
BA.2.3.7	3	0	3
BA.2.3.8	2	0	2
BA.2.31	17	0	17
BA.2.32	4	0	4
BA.2.34	2	0	2
BA.2.35	5	0	5
BA.2.36	27	0	27
BA.2.37	120	0	120
BA.2.38	704	11	715
BA.2.38.1	3	0	3
BA.2.38.2	3	0	3
BA.2.40.1	24	0	24
BA.2.41	4	0	4
BA.2.42	2	0	2
BA.2.43	1	0	1
BA.2.47	2	0	2
BA.2.48	15	0	15
BA.2.49	4	0	4
BA.2.5	6	0	6
BA.2.50	11	0	11
BA.2.51	22	0	22
BA.2.52	9	0	9
BA.2.53	1	0	1
BA.2.54	1	0	1
BA.2.55	1	0	1
BA.2.56	45	1	46
BA.2.57	2	0	2
BA.2.59	1	0	1
BA.2.6	7	0	7
BA.2.62	4	0	4
BA.2.64	14	0	14
BA.2.65	1,722	0	1,722
BA.2.66	23	0	23
BA.2.68	6	0	6
BA.2.7	16	0	16
BA.2.72	6	0	6
BA.2.73	1	0	1

WHO label / Pango lineage	January 1, 2021 – August 6, 2022	August 7 – September 3, 2022	Total
BA.2.74	41	4	45
BA.2.75	60	63	123
BA.2.75.1	4	16	20
BA.2.75.2	1	5	6
BA.2.76	87	6	93
BA.2.78	5	0	5
BA.2.79	6	0	6
BA.2.79.1	1	0	1
BA.2.8	9	0	9
BA.2.81	10	0	10
BA.2.82	1	0	1
BA.2.9	1,512	0	1,512
BA.2.9.2	2	0	2
BA.2.9.3	14	0	14
BA.2.9.4	1	0	1
BA.2.9.5	52	0	52
BA.3.1	1	0	1
BA.4	626	32	658
BA.4.1	1,202	129	1,331
BA.4.1.1	28	2	30
BA.4.1.4	3	0	3
BA.4.1.5	1	1	2
BA.4.1.6	22	3	25
BA.4.1.8	35	24	59
BA.4.2	141	23	164
BA.4.3	2	0	2
BA.4.4	76	7	83
BA.4.5	2	0	2
BA.4.6	597	396	993
BA.4.7	3	4	7
BA.5	157	73	230
BA.5.1	3,733	1,173	4,906
BA.5.1.1	603	129	732
BA.5.1.10	87	19	106
BA.5.1.2	83	32	115
BA.5.1.3	22	10	32
BA.5.1.4	9	1	10
BA.5.1.5	6	23	29
BA.5.1.6	20	9	29
BA.5.1.7	256	135	391

WHO label / Pango lineage	January 1, 2021 – August 6, 2022	August 7 – September 3, 2022	Total
BA.5.1.8	3	0	3
BA.5.1.9	0	1	1
BA.5.10	66	17	83
BA.5.2	2,673	1,667	4,340
BA.5.2.1	4,463	1,989	6,452
BA.5.2.2	68	29	97
BA.5.2.3	54	18	72
BA.5.2.4	1	9	10
BA.5.2.6	3	0	3
BA.5.3	18	0	18
BA.5.3.1	310	61	371
BA.5.3.2	9	0	9
BA.5.3.3	13	3	16
BA.5.3.4	4	0	4
BA.5.5	1,332	191	1,523
BA.5.5.1	0	11	11
BA.5.6	614	196	810
BA.5.6.1	1	0	1
BA.5.6.2	8	7	15
BA.5.8	122	18	140
BA.5.9	3	6	9
BC.2	1	0	1
BE.1	746	177	923
BE.1.1	402	109	511
BE.1.2	1	22	23
BE.2	2	3	5
BE.3	126	34	160
BF.1	221	27	248
BF.1.1	9	0	9
BF.10	848	388	1,236
BF.11	6	8	14
BF.12	11	39	50
BF.15	2	0	2
BF.2	20	12	32
BF.3	14	4	18
BF.3.1	3	0	3
BF.4	136	29	165
BF.5	231	127	358
BF.6	5	2	7
BF.7	15	42	57

WHO label / Pango lineage	January 1, 2021 – August 6, 2022	August 7 – September 3, 2022	Total
BF.8	31	25	56
BF.9	590	82	672
BG.2	85	1	86
BG.4	14	0	14
BG.5	24	0	24
BG.6	11	0	11
BH.1	1	2	3
BK.1	54	41	95
Variant of Interest (VOI)			
Mu	241	0	241
Lambda	8	0	8
Recombinant	76	0	76
XAC	29	0	29
XAF	1	0	1
XAM	5	0	5
XAN	4	0	4
XAP	3	0	3
XAQ	1	0	1
XAS	8	0	8
XB	1	0	1
XE	3	0	3
XM	1	0	1
XN	1	0	1
XQ	4	0	4
XW	14	0	14
XZ	1	0	1
Non-VOC/VOI	6,293	0	6,293
Total sequenced	155,741	7,781	163,522

Note: Results do not represent all Ontario cases. Includes results from PHO since January 1, 2021, The Hospital for Sick Children since April 21, 2021, Kingston Health Sciences Centre since January 1, 2021, Shared Hospital Laboratory since March 26, 2021, and Hamilton Regional Laboratory Medicine Program since April 11, 2021. Past testing algorithms have led to preferential sequencing of samples with N501Y and/or E484K mutations detected by PCR, which has biased the results toward lineages with these mutations. Pango lineage assignments may change over time, which may impact cumulative totals. Results should be interpreted with caution as frequencies do not reflect prevalence. Sample date represents the earliest date available for the sample. If more than one sample was sequenced for a case, the most recent sample was included.

Data sources: PHO, Hospital for Sick Children, Kingston Health Sciences Centre, Shared Hospital Laboratory, Hamilton Regional Laboratory Medicine Program

Technical Notes

Data Sources

Public Health Ontario (PHO)

- Data were extracted from the PHO Laboratory Information Management System on September 14, 2022 at approximately 2:00 a.m.
- Data were extracted from the PHO SARS-CoV-2 Whole Genome Sequencing Database on September 14, 2022 at approximately 12:00 p.m.

The Hospital for Sick Children (HSC)

- Data were received by PHO on September 13, at approximately 12:00 p.m.

Kingston Health Sciences Centre (KHSC)

- Data were received by PHO on September 13, 2022 at approximately 9:40 a.m.

Shared Hospital Laboratory (SHL)

- Data were received by PHO on September 13, 2022 at approximately 8:00 p.m.

Hamilton Regional Laboratory Medicine Program (HRLMP)

- Data were received by PHO on September 14, 2022 at approximately 12:30 a.m.

Public Health Case and Contact Management Solution (CCM)

- Data were extracted from the Public Health Case and Contact Management Solution on September 12, 2022 at approximately 1:00 p.m.

Ontario Ministry of Health's COVaxON application (COVaxON)

- COVID-19 vaccination data were based on information successfully extracted from the Ontario Ministry of Health's COVaxON application as of September 12, 2022 at approximately 7:00 a.m.
- COVaxON data was linked to COVID-19 case data from CCM.

Ontario SARS-CoV-2 Whole Genome Sequencing Strategy

- At the beginning of 2021, Ontario's whole genome sequencing strategy was to sequence samples with specific mutations identified from VOC PCR testing to confirm they were variants of concern. From February 3, 2021 this included sequencing samples with the N501Y mutation detected (initially associated with the B.1.1.7 [Alpha] lineage) and from March 22, 2021, samples with the E484K mutation detected (initially associated with the P.1 [Gamma] and B.1.351 [Beta] lineages).

- As of May 2, 2021, Ontario's strategy shifted to representative surveillance with VOC PCR testing laboratories being asked to send a proportion of eligible samples ($Ct \leq 30$ and sufficient volume remaining) to Ontario COVID-19 Genomics Network (OCGN) sequencing laboratories. PHO began sequencing a 10% systematic sample of eligible samples on May 2 and 50% on May 30. Other VOC PCR testing laboratories were asked to begin submitting a 10% systematic or random sample of eligible samples to OCGN laboratories on May 26 and 50% on June 2. All VOC PCR/diagnostic testing laboratories shifted to submitting 100% of eligible samples to the OCGN on June 14; 50% on August 27; 10% on September 10; 25% on October 8; 100% on November 5; 50% on December 7; 10% on December 20; 5% on December 30; 20% on February 16; 50% on March 9; 25% on March 30; 10% on April 13; 25% on May 12; 100% on June 10; 50% on July 8; 20% on July 22; and 50% on August 24, 2022. The proportion of samples sequenced may change over time with changes in provincial case trends. Due to logistics, not all laboratories may have implemented sampling proportion changes at the same time.
- As of November 12, 2021, VOC PCR testing has been discontinued. The 73 diagnostic testing laboratories were asked to send all eligible samples (diagnostic PCR $Ct \leq 30$ and sufficient volume remaining) to one of the five OCGN laboratories for whole genome sequencing.
- As of December 6, 2021, VOC PCR testing for S gene target failure (SGTF) was implemented across Ontario to screen for Omicron. Diagnostic testing laboratories were asked to send all eligible samples (diagnostic PCR $Ct \leq 35$ and sufficient volume remaining) to one of eight SGTF testing laboratories. The SGTF testing laboratories will then submit a proportion of eligible samples (SGTF PCR $Ct \leq 30$ and sufficient volume remaining) for WGS according to the representative surveillance strategy.
 - Due to logistics, not all laboratories may have implemented SGTF testing at the same time.
- As of December 30, 2021, SGTF testing of all eligible samples was discontinued in Ontario. The 73 diagnostic testing laboratories were asked to send a proportion of eligible samples (diagnostic PCR $Ct \leq 30$ and sufficient volume remaining) to one of the five OCGN laboratories for whole genome sequencing, according to the representative surveillance strategy.
- As of December 31, 2021, diagnostic PCR testing was restricted to high-risk populations. Since then, the Ministry of Health continues to update its [guidance on testing](#). As such, representative surveillance only pertains to tested populations.

Data Caveats and Methods: Ontario COVID-19 Genomics Network (OCGN)

- Lineage nomenclature is dynamic. Pango lineage naming and assignment may change as more samples are sequenced and analyzed globally. Similarly, VOC and VOI classifications may change.
- Whole genome sequencing sample logistics are complex and require samples to be transferred across a large network of laboratories. Samples are initially sent to one of 73 diagnostic testing laboratories. If the diagnostic PCR cycle threshold is ≤ 35 and there is sufficient volume remaining, samples are submitted for testing at one of 11 VOC PCR testing laboratories. If the VOC PCR cycle threshold is ≤ 30 and there is sufficient volume remaining, VOC PCR testing laboratories have been asked to submit a proportion of their eligible samples to one of five

OCGN laboratories for sequencing according to the surveillance strategy. As of November 12, VOC PCR has been discontinued. Diagnostic testing laboratories now send eligible samples ($Ct \leq 30$ and sufficient volume remaining) directly to one of the five OCGN laboratories for whole genome sequencing.

- PHO is unable to confirm whether VOC PCR testing laboratories or diagnostic testing laboratories (November 12, 2021 onwards) have submitted eligible samples.
- Data submitted to PHO from OCGN laboratories have not been independently verified.
- The dates associated with samples submitted by network laboratories vary due to sample logistics and different laboratory information systems. Dates associated with WGS samples were assigned based on a hierarchy: sample collection date > SARS-CoV-2 diagnostic received date > SARS-CoV-2 diagnostic reported date > VOC PCR received date > VOC PCR reported date > WGS received date > WGS reported date. Weeks were created to align with surveillance weeks used by the Public Health Agency of Canada for influenza reporting.
- Samples from the same case were linked if they had the same health card number or if they had the same first name, last name, and date of birth. If more than one sample was sequenced for a case, the most recent sample was used. This may shift a case to a more recent week if a subsequent sample was sequenced from the same case. A small proportion of cases may have samples that were not linked due to inconsistencies or data entry errors.
- Results for recent weeks are incomplete as not all sequencing and bioinformatics analyses were complete at the time of data extraction.
- Public health unit was assigned using diagnosing health unit in CCM. If the case did not link to CCM (2.9%), then public health unit was assigned using OCGN patient postal code or ordering provider postal code if patient postal code was missing.
- For representative surveillance, results may not be representative of Ontario overall. Samples selected include a proportion of eligible samples received by OCGN laboratories according to the whole genome sequencing strategy. Individual VOC PCR laboratories may have implemented the strategy and/or increased the proportion of samples selected on different dates. Cumulative data included do not reflect all whole genome sequencing conducted in Ontario.
 - Data from the OCGN laboratories cover different time periods: PHO since January 1, 2021, HSC since April 21, 2021, KHSC since January 1, 2021, SHL since March 26, 2021, and HRLMP since April 11, 2021.
 - Past testing algorithms have led to preferential sequencing of samples with N501Y and/or E484K mutations detected by VOC PCR. This has created a sampling bias reflected in the distribution of lineage results prior to May 2, 2021.

Methods: Nowcast Estimates, Projections and Weekly Relative Growth Rate

- Nowcast estimates and projections are generated using a multinomial logistic regression model. The Nowcast model uses twelve weeks of daily representative surveillance data up to the most recent date, with date as the univariate model predictor. Lineages that had at least one day with an estimated or projected prevalence of 5% or greater were included in the table and figure. Lineages with at least seven days of non-zero case counts were included.
- Projected Nowcast estimates are future predictions of prevalence after the most recent date of observed data.
- Relative growth rate is a measure of a lineage's growth relative to a reference lineage.⁶ Relative growth rates greater than 1 suggest an increased growth rate compared to the reference; relative growth rates less than 1 suggest a decreased growth rate compared to the reference.
- These relativized growth rates can be calculated by exponentiating the selection rate coefficient from the Nowcast model.
 - The selection rate coefficient is the difference in growth rate between two lineages ($\Delta r = r_{\text{lineage}} - r_{\text{reference}}$), and can be derived from a logistic regression model where the outcome is the relative frequency of a lineage and the predictor is time.^{6,7}

Data Caveats and Methods: Public Health Case and Contact Management Solution (CCM)

- CCM is a dynamic disease reporting system, which allows ongoing updates to data previously entered. As a result, data extracted from CCM represent a snapshot at the time of extraction and may differ from previous or subsequent reports.
- Methods for processing the CCM case data are described in the [Technical Notes](#) of the COVID-19 Data Tool.
- Data corrections or updates can result in case records being removed and/or updated from past reports.
- Dates associated with COVID-19 cases in Ontario were assigned using a hierarchy to best align with the sample date used for representative surveillance: sample collection date > test reported date > case reported date. As a result, the number of cases may differ from other reports which use different dates.
- Cases were linked to CCM if they had the same health card number or if they had the same first name, last name, and date of birth. Cases may not have linked to CCM due to discrepancies in patient identifiers or if they were not residents of Ontario (diagnosing health unit was reported as MOH).
- Tables for outbreak, hospitalized, and deceased indicators only include cases that linked to CCM (97.1% of cases).

- ‘Outbreak-associated cases’ include cases linked to a confirmed outbreak as declared by the local medical officer of health or their designate in accordance to the Health Protection and Promotion Act and criteria outlined in [Ministry guidance documents](#).
- Outbreaks in settings outside of Ontario are excluded from all outbreak counts.
- Data on hospital admissions and deaths are likely under-reported as these events may occur after the completion of public health follow up of cases. Cases that were admitted to hospital or died after follow-up was completed may not be captured in CCM.
- Hospital admission includes all cases ever admitted to a hospital (or that had their hospital stay extended) because of COVID-19. It includes cases that have been discharged from hospital as well as cases that are currently hospitalized. Includes cases in an Intensive Care Unit (ICU) but not emergency room visits. Hospital admissions were identified by a reported hospital admission date or reported ‘Yes’ for hospitalization/ICU.
- For surveillance purposes, a COVID-19 death is defined as a death resulting from a clinically compatible illness unless there is a clear alternative cause of death that cannot be related to COVID-19 (e.g., trauma, medically assisted death). There should be no period of complete recovery from COVID-19 between illness and reported death.
- Deaths are determined by using the outcome and Type of Death fields in CCM. COVID-19 deaths are counted where the Outcome value is ‘Fatal’ and the Type of Death value is not ‘DOPHS was unrelated to cause of death’ or ‘Under PHU Review’.

Data Caveats and Methods: COVaxON

- In order to identify cases post-vaccination, vaccine uptake data extracted from the Ontario Ministry of Health’s (MOH) COVaxON application was linked to case data extracted from the MOH’s Public Health Case and Contact Management Solution (CCM).
 - Clients in COVaxON and CCM were linked using health card number as well as other personal identifiers, including name, date of birth, gender, and postal code.
- Linkage was done using processed COVaxON and CCM data. Methods for processing COVaxON vaccine uptake data are described in the Technical Notes of the [COVID-19 Vaccine Uptake Report](#) and methods for processing post-vaccination cases are described in the Technical Notes of the [Confirmed Cases of COVID-19 Post Vaccination Report](#).
- Only cases that have received Health Canada authorized vaccines including, Pfizer-BioNTech Comirnaty™, Moderna Spikevax™, AstraZeneca Vaxzevria™/COVISHIELD, and Janssen are included. Cases that received one or more doses of a non-Health Canada authorized vaccine are excluded.
- A higher proportion of cases reported in post-series initiation and post-series completion is a reflection of both trends in vaccine administration (increasing number of doses administered over time) and trends in COVID-19 incidence. Further details on vaccine administration trends in Ontario are described in the [COVID-19 Vaccine Uptake Report](#).

- Factors, such as age, that may affect the risk of COVID-19 hospitalization and death are not accounted for in these analyses. As a result, the proportion of severe outcomes following booster doses may be higher compared to other vaccination statuses since booster eligibility was initially restricted to older adults who are more likely to have severe outcomes and who also have the highest uptake for booster doses.
- For vaccine category definitions, please refer to [Confirmed Cases of COVID-19 Following Vaccination in Ontario](#).

References

1. Public Health Agency of Canada. SARS-CoV-2 variants: national definitions, classifications and public health actions [Internet]. Ottawa, ON: Government of Canada; 2021 [modified 2021 August 26; cited 2021 August 27]. Available from: <https://www.canada.ca/en/public-health/services/diseases/2019-novel-coronavirus-infection/health-professionals/testing-diagnosing-case-reporting/sars-cov-2-variants-national-definitions-classifications-public-health-actions.html>
2. World Health Organization. Tracking SARS-CoV-2 variants [Internet]. Geneva, Switzerland; 2021 [modified 2021 October 29; cited 2021 November 1]. Available from: <https://www.who.int/en/activities/tracking-SARS-CoV-2-variants/>
3. European Centre for Disease Prevention and Control. SARS-CoV-2 variants of concern as of 28 October 2021 [Internet]. Stockholm, Sweden; 2021 [modified 2021 October 28; cited 2021 November 1]. Available from: <https://www.ecdc.europa.eu/en/covid-19/variants-concern>
4. Rambaut A, Holmes EC, O'Toole Á, Hill V, McCrone JT, Ruis C, et al. A dynamic nomenclature proposal for SARS-CoV-2 lineages to assist genomic epidemiology. *Nat Microbiol*. 2020; 5(11):1403-7. Available from: <https://doi.org/10.1038/s41564-020-0770-5>
5. cov-lineages. pangolin [Internet]. 2020 [cited 2021 May 29]. GitHub. Available from: <https://github.com/cov-lineages/pangolin>
6. Davies N, Abbott S, Barnard R, Jarvis CI, Kucharski AJ, Munday JD, et al. Estimated transmissibility and impact of SARS-CoV-2 lineage B.1.1.7 in England. *Science*. 2021; 372(6538): eabg3055. Available from: <https://doi.org/10.1126/science.abg3055>
7. Campbell F, Archer B, Laurenson-Schafer H, Jinnai Y, Konings F, Batra N, et al. Increased transmissibility and global spread of SARS-CoV-2 variants of concern as at June 2021. *Euro Surveill*. 2021; 26(24). Available from: <https://doi.org/10.2807/1560-7917.ES.2021.26.24.2100509>

Citation

Ontario Agency for Health Protection and Promotion (Public Health Ontario). Epidemiologic summary: SARS-CoV-2 whole genome sequencing in Ontario, September 16, 2022. Toronto, ON: King's Printer for Ontario; 2022.

Disclaimer

This document was developed by Public Health Ontario (PHO). PHO provides scientific and technical advice to Ontario's government, public health organizations and health care providers. PHO's work is guided by the current best available evidence at the time of publication. The application and use of this document is the responsibility of the user. PHO assumes no liability resulting from any such application or use. This document may be reproduced without permission for non-commercial purposes only and provided that appropriate credit is given to PHO. No changes and/or modifications may be made to this document without express written permission from PHO.

For Further Information

For more information, email communications@oahpp.ca.

Public Health Ontario

Public Health Ontario is an agency of the Government of Ontario dedicated to protecting and promoting the health of all Ontarians and reducing inequities in health. Public Health Ontario links public health practitioners, front-line health workers and researchers to the best scientific intelligence and knowledge from around the world.

For more information about PHO, visit publichealthontario.ca.